

Original Paper

Assessment of Physical Activity by Wearable Technology During Rehabilitation After Cardiac Surgery: Explorative Prospective Monocentric Observational Cohort Study

Isabeau Thijs^{1*}, MSc; Libera Fresiello^{2,3*}, MSc, PhD; Wouter Oosterlinck¹, MD, PhD; Peter Sinnaeve⁴, MD, PhD; Filip Rega¹, MD, PhD

¹Research Unit of Cardiac Surgery, Department of Cardiovascular Sciences, University Hospitals Leuven, Leuven, Belgium

²Department of Cardiac Surgery, Katholiek Universiteit Leuven, Leuven, Belgium

³Institute of Clinical Physiology, National Research Council, Pisa, Italy

⁴Research Unit of Cardiology, Department of Cardiovascular Sciences, University Hospitals Leuven, Leuven, Belgium

*these authors contributed equally

Corresponding Author:

Isabeau Thijs, MSc

Research Unit of Cardiac Surgery

Department of Cardiovascular Sciences

University Hospitals Leuven

Herestraat 49, Bus 7003

Leuven, 3000

Belgium

Phone: 32 +3216344219

Email: isabeau.thijs@outlook.com

Abstract

Background: Wearable technology is finding its way into clinical practice. Physical activity describes patients' functional status after cardiac surgery and can be monitored remotely by using dedicated trackers.

Objective: The aim of this study was to compare the progress of physical activity in cardiac rehabilitation by using wearable fitness trackers in patients undergoing coronary artery bypass surgery by either the conventional off-pump coronary artery bypass (OPCAB) or the robotically assisted minimally invasive coronary artery bypass (RA-MIDCAB). We hypothesized faster recovery of physical activity after RA-MIDCAB in the first weeks after discharge as compared to OPCAB.

Methods: Patients undergoing RA-MIDCAB or OPCAB were included in the study. Each patient received a Fitbit Charge HR (Fitbit Inc, San Francisco, CA) physical activity tracker following discharge. Rehabilitation progress was assessed by measuring the number of steps and physical activity level daily. The physical activity level was calculated as energy expenditure divided by the basic metabolic rate.

Results: A total of 10 RA-MIDCAB patients with a median age of 68 (min, 55; max, 83) years and 12 OPCAB patients with a median age of 69 (min, 50; max, 82) years were included. Baseline characteristics were comparable except for body mass index (RA-MIDCAB: 26 kg/m²; min, 22; max, 28 versus OPCAB: 29 kg/m²; min, 27; max, 33; $P < .001$). Intubation time ($P < .05$) was significantly lower in the RA-MIDCAB group. A clear trend, although not statistically significant, was observed towards a higher number of steps in RA-MIDCAB patients in the first week following discharge.

Conclusions: RA-MIDCAB patients have an advantage in recovery in the first weeks of revalidation, which is reflected by the number of steps and physical activity level measured by the Fitbit Charge HR, as compared to OPCAB patients. However, unsupervised assessment of daily physical activity varied widely and could have consequences with regard to the use of these trackers as research tools.

(JMIR Mhealth Uhealth 2019;7(1):e9865) doi:[10.2196/mhealth.9865](https://doi.org/10.2196/mhealth.9865)

KEYWORDS

fitness trackers; coronary artery bypass; cardiac surgery; cardiac rehabilitation; postoperative care; wearable; physical activity; exercise

Introduction

Kolesov V (1964) performed the first coronary artery bypass graft (CABG) using the internal mammary artery to treat a patient with ischemic myocardial heart disease [1]. Off-pump coronary artery bypass (OPCAB) surgery was developed to reduce potential adverse effects induced by the use of cardiopulmonary bypass and cardioplegic arrest [2]. In this approach, the harvesting of the internal mammary arteries and anastomoses are performed on a beating heart through a median sternotomy.

Robotically assisted minimally invasive direct coronary artery bypass (RA-MIDCAB) aims to further reduce the invasiveness of the OPCAB approach by avoiding midline sternotomy. In RA-MIDCAB surgery, the internal mammary arteries are prelevated via a thoracoscopy using robotic assistance. The grafting of the bypass is performed in a second stage via a small (4-5 cm) left anterolateral minithoracotomy. Similar to OPCAB, the anastomosis is performed on a beating heart, without the use of cardiopulmonary bypass. The RA-MIDCAB approach reduces morbidity, length of hospital stay, need for blood transfusion, and wound infections [3-5]. The time for recovery is controversial among studies [3-5]: Some indicate an earlier recovery to full physical activity [3,6].

After cardiac surgery, patients need structured support to improve functional capacity and restore their quality of life. Phase II cardiac rehabilitation programs are developed to deliver comprehensive support such as monitoring in physical and psychological conditions and education of patients on healthy long-term routines. Phase II cardiac rehabilitation is suggested as a class I recommendation in the treatment of cardiac diseases by the European Society of Cardiology, the American Heart Association, and the American College of Cardiology [7-9]. Adherence could potentially be tracked by the use of remote monitoring systems.

Physical activity or fitness trackers are wearable sensors, often worn as a wristband or embedded in a smartwatch or mobile phone, that collect data on one's daily physical activity. Most of these commercially available trackers include an accelerometer to assess step counts; distance walked; and intensity, duration, and type of movement (eg, walking, running, and jogging). Thus, users can have direct access to their personal data and take an active role in monitoring their health [10,11].

These trackers are also of use in clinical practices and research. Accelerometry data can be derived noninvasively and in unsupervised, free-living conditions, which provides an opportunity to better describe patients' activity of daily living and health status in terms of mobility, behavioral pattern, and functional ability. Consequently, these data can contribute to more comprehensive, relevant, and high-quality clinical research data [10]. In clinical practices, home telemonitoring trials show favorable results in pulmonary and cardiac patients [12]. In

cardiac rehabilitation, multiple cardiac telecare trials have shown a noninferiority or superiority of telemonitoring and telecoaching of patients in a cardiac rehabilitation program compared to conventional center-based supervised cardiac rehabilitation programs [12,13,14]. These physical activity trackers have the ability to encourage exercise and lifestyle behavior and monitor and share progress [11,12,15]. As such, wearables could potentially have a future in at-home management and remote monitoring of patients with chronic diseases and in secondary preventive care after cardiac surgery.

At the University Hospital of Leuven, an explorative clinical observational study was performed to evaluate physical activity in patients after coronary artery bypass (CAB) surgery. This study aimed to quantify physical fitness at particular time points and investigate whether minimal-access surgical procedures can assure faster recovery and better outcome than the conventional, more invasive surgical procedure.

Methods

Study Design

Protocol

An explorative prospective monocentric observational cohort study was conducted at the Cardiac Surgery Department of the University Hospitals of Leuven. The clinical protocol conformed to the principles outlined in the Declaration of Helsinki and was approved by the ethical committee of the University Hospitals Leuven. All patients provided written informed consent prior to inclusion in the study.


Patients

The study included patients with coronary artery disease who were eligible for elective surgical revascularization according to the most recent guidelines of the European Society of Cardiology [16]. They were scheduled to undergo either an RA-MIDCAB or OPCAB procedure. In both approaches, grafts are anastomosed on the diseased vessels without the use of cardiopulmonary bypass—the so-called off-pump technique. In OPCAB, a full sternotomy is performed, whereas in RA-MIDCAB, the anastomosis is performed through a small left anterolateral thoracotomy. Additionally, in RA-MIDCAB, internal mammary arteries (used as grafts) are harvested using robotic assistance from the Da Vinci Surgical System (Intuitive Surgical Inc, Sunnyvale, CA).

Eligibility Criteria

The main exclusion criteria were urgently scheduled and on-pump procedures, mobility problems that could interfere with physical activity, and the presence of cognitive impairment that prevented subjects from fully understanding the protocol. An overview of the inclusion and exclusion criteria is provided in [Multimedia Appendix 1](#).

Figure 1. Timeline of the study design constructed in three evaluation time points: a preoperative inclusion, a 14-day Fitbit-wearing period after discharge, and a 7-day Fitbit-wearing period after a follow-up consult, 4 weeks after discharge. ICF: Informed Consent Form; DAT: Daily Activity Tracking.



Evaluation Design

The protocol was organized in three evaluation time points: a preoperative baseline assessment and two periods of wearing a physical activity tracker (Figure 1). At discharge, patients received a Fitbit Charge HR (Fitbit Inc, San Francisco, CA) and were asked to wear the wearable device for 14 consecutive days. Four weeks later, a follow-up consultation was scheduled, and the patient was again asked to wear the device for 7 days. The patients were asked to return the tracker to the hospital by mail. Additionally, throughout the hospital stay, clinical data were collected. An overview of the variables for which data were collected is shown in [Multimedia Appendix 2](#).

Physical Fitness Assessment

Subjects' daily physical activity was described by parameters recorded by the Fitbit Charge HR. This activity tracker is a wrist band with an interface through which patients can monitor their real-time progress. The tracker was set according to the height, weight, and age of the subject. Subjects were instructed to wear the wristband as much as possible during the day. The tracker records the daily number of steps, the distance walked, the flights of stairs taken, the intensity and duration of exercise, the estimated energy expenditure, the sleeping pattern and the heart rate variation using the Pure Pulse technology (FitBit Inc).

Data Analysis

Data from the Fitbit Charge HR were analyzed by calculating the weekly average step counts and energy expenditure (kcal). For energy expenditure, the physical activity level was calculated by dividing the total caloric expenditure by the basic metabolic rate. This physical activity level represents the physical activity adjusted for weight, height, and age (included in the basic metabolic rate). For every subject, the first (discharge) and last (return) day of Fitbit wearing were excluded, since biased results were expected. Furthermore, continuous heart rate was evaluated to check periods when patients did not wear the device. Up to 2 hours a day of non-wearing time were neglected; if the non-wearing period was longer, that day was excluded. This time loss could be due to battery charging or activities such as bathing or showering.

Outcome Measures

Primary Outcome

The primary outcome was the objective physical activity score described by the Fitbit activity tracker data during the two

periods mentioned above in Evaluation Design. Weekly average number of steps and weekly average physical activity level were used to quantify physical activity.

Secondary Outcomes

Secondary outcomes included observational data including demography, cardiac and noncardiac history, operative variables, and postoperative complications until 4 weeks after discharge.

Statistical Analyses

Statistical analyses were carried out using IBM SPSS Statistics, Version 24 (IBM Corp, Armonk, NY). Since only a small sample size was included, differences in continuous variables were analyzed using two-sided Mann Whitney *U* tests and reported as median and range with minimal and maximal values. For dichotomous variables, a Fisher exact test was performed, and for categorical variables, a Mann Whitney *U* test was performed. For repeated measures analysis, the nonparametric Friedman *t* test was used. Statistical significance was considered for *P* values < .05.

Results

Patient Recruitment

Patients were recruited from January 2017 to April 2017. In total, 25 patients were enrolled, of which 11 were RA-MIDCAB patients and 14 were OPCAB patients. Three patients were excluded after the surgery: one patient was excluded from the RA-MIDCAB group due to a prolonged hospital stay as a consequence of acute on chronic kidney failure, and two patients dropped out after surgery in the OPCAB group (one withdrew from the study and one died). Furthermore, after the 14-day Fitbit-wearing period, three patients in the OPCAB group and two patients in the RA-MIDCAB group dropped out. A study flowchart is presented in [Figure 2](#). Results of the baseline characteristics are listed in [Table 1](#).

The two groups did not significantly differ in age ($P=.79$), gender ($P=>.99$), height ($P=.79$), and weight ($P=.07$), but OPCAB patients had a significantly higher body mass index (29 kg/m^2) than RA-MIDCAB patients (26 kg/m^2 ; $P<.001$).

Heart failure distribution according to the New York Heart Association class was not significantly different in both groups ($P=.89$), and most subjects belonged to class I and II. The median left ventricular ejection fraction was 60% (min, 45; max, 78) in the RA-MIDCAB group and 60% (min, 40; max, 78) in the OPCAB group. Furthermore, both groups showed a similar

distribution in the European System for Cardiac Operative Risk Evaluation (EuroSCORE II; $P=.21$), left ventricular ejection fraction ($P=.69$), history of arrhythmias ($P=.46$), history of myocardial infarction ($P=.90$), and mitral regurgitation ($P>.99$).

There was a trend towards the presence of hypercholesterolemia in OPCAB patients compared to RA-MIDCAB patients (92% and 50%, respectively, $P=.06$). No significant differences were observed in the presence of comorbidities (Table 1).

Intubation time was significantly higher in the OPCAB group ($P<.05$), with 15 hours 30 minutes (min, 8 hours 2 minutes; max, 21 hours 50 minutes) in contrast to 8 hours 45 minutes (min, 5 hours 49 minutes; max, 23 hours) in the RA-MIDCAB group (Table 1). In addition, the number of grafts was significantly higher in the OPCAB group ($P<.005$), and the operation duration was higher in the OPCAB group ($P=.10$).

The median duration was 5 hours 15 minutes (min, 3 hours 7 minutes; max, 6 hours 58 minutes), whereas the median duration in the RA-MIDCAB group was 4 hours 40 minutes (min, 3 hours 18 minutes; max, 5 hours 26 min).

Postoperatively, five patients in the RA-MIDCAB group and one patient in the OPCAB group were transferred to the postanesthetic care unit; a trend towards significance was observed in this parameter ($P=.06$). The length of stay at the postoperative care units (including the postanesthetic care unit and intensive care unit) was significantly lower in the RA-MIDCAB group ($P<.001$), with a median stay of 20 hours 45 minutes (min, 15 hours 30 minutes; max, 45 hours) in this group and 30 hours 45 minutes (min, 18 hours 30 minutes; max, 77 hours) in the OPCAB group. Furthermore, the overall hospital stay was similar in both groups ($P=.21$).

Figure 2. Study flowchart. CABG: coronary artery bypass graft; RA-MIDCAB: robotically assisted minimally invasive direct coronary artery bypass; OPCAB: off-pump coronary artery bypass; AKI: acute kidney injury.




Table 1. Baseline characteristics and postoperative parameters.

Characteristics	RA-MIDCAB ^a (n=10)	OPCAB ^b (n=12)	P value
Demographics			
Sex (male), n (%)	9 (90)	10 (78)	>.99
Age (y), median (min, max)	68 (55, 83)	69 (50, 82)	.79
Height (cm), median (min, max)	172 (154, 178)	171 (155, 178)	.79
Weight (kg), median (min, max)	77 (57, 90)	83 (75, 100)	.07
Body mass index (kg/m ²), median (min, max)	26 (22, 28)	29 (27, 33)	<.005 ^c
Cardiac history			
NYHA^d Class, n			.89
I	4	5	
II	4	5	
III	2	2	
Presence of unstable angina, n	2	3	.79
Left ventricular ejection fraction (%), median (min, max)	60 (45, 78)	60 (40, 78)	.69
EuroSCORE II ^e (%), median (min, max)	1.7 (0.6, 5.5)	1.1 (0.6, 2.8)	.21
History of arrhythmias, n	1	0	.46
History of myocardial infarction, n			.9
Non-ST-segment elevated myocardial infarction, n	2	3	
ST-segment elevated myocardial infarction, n	1	0	
Mild mitral regurgitation, n	5	5	>.99
Non-cardiac history			
Smoking status, n			.64
Ex-smoker since >1 month	5	6	
Smoker	3	1	
Diabetes mellitus type II, n	3	2	.62
Arterial hypertension, n	7	10	.62
Pulmonary hypertension, n ^f	1	1	>.99
Hypercholesterolemia, n	5	11	.06
Renal impairment, n	9	6	.83
Peripheral vascular diseases, n	1	2	>.99
Chronic obstructive pulmonary disease, n	1	2	>.99
Operative data, median (min, max)			
Operation duration, min	281 (198, 326)	316 (187, 418)	.1
Intubation time, min	527 (349, 1380)	931 (482, 1310)	<.05 ^c
Number of grafts, n	2 (1, 2)	3.5 (2, 4)	<.005 ^c
Postoperative data			
Postanesthetic care unit stay, n	5	1	
Postoperative care unit length of stay (min), median (min, max)	1245 (930, 2700)	1440 (1110, 4620)	<.001 ^c
Hospital length of stay (d), median (min, max)	6 (4, 12)	7 (5, 15)	.21
Complications, n			
Wound infections	0	2	.48

Characteristics	RA-MIDCAB ^a (n=10)	OPCAB ^b (n=12)	P value
Pulmonary infections	0	2	.48
Pleural effusions	2	3	>.99
New arrhythmias	2	1	.57
Hypokalemia	4	3	.65
Pericarditis	2	0	.2

^aRA-MIDCAB: robotically assisted minimally invasive direct coronary artery bypass.

^bOPCAB: off-pump coronary artery bypass.

^cP values are significant.

^dNYHA: New York Heart Association functional classification of heart failure.

^eEuroSCORE II: European System for Cardiac Operative Risk Evaluation.

^fPulmonary hypertension was defined as pulmonary pressure >25 mmHg.

Table 2. Step count analysis recorded by Fitbit Charge HR in RA-MIDCAB and OPCAB patients.

Average step counts	RA-MIDCAB ^a	Percentage ^b	OPCAB ^c	P value
Overall analysis, median (min, max), n				
Week 1	3715 (1637, 6720), 10	335	1110 (739, 10,195), 11	.06
Week 2	4357 (1415, 7671), 10	238	1832 (856, 11,282), 10	.33
Week 5	6012 (3473, 11579), 8	105	5719 (2128, 11,948), 9	.7
Analysis without dropouts, median (min, max), n				
Week 1	3715 (1734, 6720), 8	371	1001 (739, 10,195), 9	.07
Week 2	4357 (1512, 7286), 8	459	949 (856, 11,282), 9	.17
Week 5	6012 (3473, 11579), 8	105	5719 (2128, 11,948), 9	.7
Repeated measures Friedman <i>t</i> test				
Chi-square	28		30	— ^d
P value	<.001		<.001	—

^aRA-MIDCAB: robotically assisted minimally invasive direct coronary artery bypass.

^bPercentage of number of steps in RA-MIDCAB patients compared to OPCAB patients.

^cOPCAB: off-pump coronary artery bypass.

^dNot applicable.

Primary Outcome

Step counts

Data were corrected for periods when the physical activity tracker was not worn. In week 1, a total of 3 days for one patient were excluded from the analysis. In week 2, a total of 9 days, distributed over four patients, were excluded. In week 5, 2 days, distributed over two patients, were excluded.

In the first week, the RA-MIDCAB group showed a higher average number of steps than the OPCAB group, a result almost statistically significant ($P=.06$). Similarly, in the second week after discharge, RA-MIDCAB patients took more steps, but no significant difference was observed between the groups ($P=.33$). In week 5, the OPCAB group bridged the gap in the number of

steps, and the average number of steps was similar between the two groups ($P=.70$; Table 2; Figures 3 and 4).

A nonparametric Friedman *t* test was performed to analyze the repeated measures analysis for the number of steps over time. A significant change over time was observed in the RA-MIDCAB group (28 steps; $P<.001$) and the OPCAB group (30 steps; $P<.001$; Table 2).

Physical Activity Level

With regard to the physical activity level, no significant differences were observed in weeks 1, 2, and 5 between the RA-MIDCAB and OPCAB groups ($P=.36$, $P=.36$, and $P=.50$, respectively). However, the physical activity level was higher in the RA-MIDCAB group than in the OPCAB group in all weeks (Table 3).

Figure 3. Weekly average number of steps in robotically assisted minimally invasive direct coronary artery bypass (RA-MIDCAB) and off-pump coronary artery bypass (OPCAB) patients plotted over time. Weekly average step count is plotted as median over time. n indicates the number of patients included in the cohort result. P value is for the Mann Whitney U test for the difference between the two groups at that time point. χ^2 results of repeated measures Friedman t test.




Figure 4. Boxplots of weekly average number of steps in robotically assisted minimally invasive direct coronary artery bypass (RA-MIDCAB) and off-pump coronary artery bypass (OPCAB) patients. Weekly average step counts is shown as box and whisker plots, presenting medians, 25% and 75% quartiles, minimums, and maximums. n indicates the number of patients included in the cohort result. M_W1: MIDCAB result in week 1; O_W1: OPCAB results in week 1; M_W2: MIDCAB results in week 2; O_W2: OPCAB results in week 2; M_W5: MIDCAB results in week 5; O_W5: OPCAB results in week 5.




Table 3. Physical activity level^a analysis recorded by Fitbit Charge HR in RA-MIDCAB and OPCAB patients.

Time point	RA-MIDCAB ^b		OPCAB ^c		P value
	median (min, max)	n	median (min, max)	n	
Overall analysis					
Week 1	1.39 (1.05, 1.71)	10	1.29 (1.08, 1.59)	11	.36
Week 2	1.41 (1.04, 1.63)	10	1.32 (1.04, 1.60)	10	.36
Week 5	1.52 (1.13, 1.90)	8	1.44 (1.16, 1.80)	9	.5
Analysis without dropouts					
Week 1	1.39 (1.05, 1.71)	8	1.26 (1.08, 1.59)	9	.4
Week 2	1.41 (1.04, 1.63)	8	1.23 (1.04, 1.60)	9	.42
Week 5	1.52 (1.13, 1.90)	8	1.44 (1.16, 1.80)	9	.5

^aPhysical activity level calculated as total energy expenditure divided by basic metabolic rate.

^bRA-MIDCAB: robotically assisted minimally invasive direct coronary artery bypass.

^cOPCAB: off-pump coronary artery bypass; data are reported as median (min, max).

Discussion

Principal Findings

We evaluated physical activity in cardiac rehabilitation by using the Fitbit Charge HR tracker device after conventional and minimally invasive CAB surgery. A clear trend was observed towards a higher physical activity level in RA-MIDCAB patients

than in OPCAB patients, which was reflected in the number of steps and physical activity level, although statistical significance was not reached.

Value of Wearable Activity Trackers in Surgical Outcome Research

To our knowledge, this is the first study to use wearable activity tracking in a clinical environment to compare the outcome of

two types of cardiac surgery interventions. The Fitbit Charge HR provided useful information about patients' physical activity in this study. Wearable activity trackers are finding their way in research and medical practices [10,15,17]. An important limitation, however, is that commercially available activity trackers are often not thoroughly validated for their accuracy and reliability. Studies showed that step count accuracy is dependent on gait patterns in healthy volunteers [18,19]. In the elderly and chronically ill, a negative correlation was found between the gait pattern and step count accuracy, as assessed by other commercially available activity trackers [18,20]. Postsurgical patients who are still in recovery and probably walk at a slow speed will create a bias in the number of steps counted. In addition, the Fitbit Charge HR is still rarely used in research. However, older-generation models have been tested for their step count accuracy [21-24] and energy expenditure estimation [25,26].

Taking into account the limitations of the Fitbit technology, it is worth highlighting that our data were not analyzed as exact results but were only used to compare the RA-MIDCAB and OPCAB groups. Any error in the step count technology would affect RA-MIDCAB and OPCAB patients in a similar way without impacting the comparative analysis performed in this work. No baseline references are available, and Fitbit does not disclose algorithms or mean error values. Hence, it was not possible to apply mean error corrections. However, it is distinctively true that further research is needed before commercially available self-monitoring wearables can be used in clinical applications.

Besides the lack of validation, wearable activity trackers have a promising future. Activity trackers provide the possibility for patients to monitor their activity patterns and share their progress with physicians, friends, and family members. Therefore, these trackers can be used as motivational tools to reach and maintain a healthy active lifestyle [15]. In this study, subjects were nearly always compliant and motivated to wear the tracker and monitor their own progress. Physical activity is one of the most health-enhancing practices, especially in primary and secondary prevention of cardiovascular risk factors. Physical activity counselling (by use of wearable activity monitors) has been shown to improve healthy lifestyles [27-29]. Savage et al (2008) found significant correlations between the daily number of steps in the first weeks of Phase II cardiac rehabilitation and cardiovascular risk factors [30].

The future for this wearable activity trackers is still unknown, but its implementation in medical practice would provide many benefits, for instance, in cardiac rehabilitation to overcome barriers to cardiac rehabilitation programs. Multiple trials have shown that wearable activity tracking and digital health devices encourage patients to improve their physical behavior and are therefore useful tools in cardiac rehabilitation. The Telerehab III trial showed that telerehabilitation, through use of a commercially available accelerometer, provided substantial and persistent health benefits and novel, cost-efficient care [31].

Comparison of Minimally Invasive and Conventional Coronary Artery Bypass

Our study showed that both groups were comparable. No significant differences were found in baseline characteristics, except for body mass index and the number of anastomosed grafts. EuroSCORE II calculation showed no significant difference in the predicted operative mortality. OPCAB patients have a more pronounced disease and may be considered more unwell, which can be reflected in the body mass index. In this group of patients, however, it seems that body mass index does not have a significant influence on physical activity and performance. Spearman correlation analysis showed that weight/body mass index was not related with the number of steps at any time point (Multimedia Appendix 3). Surgical and postsurgical data reflect the difference between both procedures. RA-MIDCAB entails a shorter intubation time and a shorter stay at postoperative care units, with more patients transferred to the postanesthetic care unit postoperatively (fast-track treatment). These findings are in line with previous studies [4,32]. Hospital length of stay, however, showed no significant difference between the two study groups; this is in contrast to the findings of other studies [33-35], which could be due to the small sample size of the present study. However, this variable might be dependent on institutional protocols and decision making of physicians and surgeons [36].

Despite the similar length of stay and baseline characteristics between the two groups, a clear trend was observed toward higher physical activity reflected in step counts and physical activity level in RA-MIDCAB patients in the first weeks, although statistical significance was not reached. Step counts depict the actual daily walking of patients during the day. The physical activity level depicts the energy expenditure as a result of activity, adjusted for individual basic metabolic rate. Therefore, both parameters interpret physical activity in a different manner and should be interpreted accordingly. It is harder to reach significance in the physical activity level analysis due to the smaller scale of variations. Owing to its explorative nature, this study is probably underpowered to detect smaller differences and is influenced greatly by outliers. Both groups showed significant changes in the number of steps over time (Table 2). Together with the differences between the groups, this could indicate that RA-MIDCAB patients advance in the early stages and OPCAB patients need some time to catch up. It must be noted that physical activity varied greatly among subjects, which could be due to the accuracy levels and algorithms of the device itself. In addition, physical activity is dependent on personal habits and character, referred to as self-efficacy [37,38], and the motivational support from the environment (relatives and friends). Patients who are sedentary before the surgery would likely abide by this lifestyle after surgery. Patients who are regularly active would probably be more motivated to achieve their prior level of fitness before the disease became symptomatic. This was illustrated in the Telerehab III trial where patients partially relapsed after telerehabilitation was stopped [31]. However, this effect would influence both groups similarly. Furthermore, inclusion in a clinical study and the intervention for monitoring activity by use of a tracker could be motivating factors. The tracker makes

it possible for patients to monitor their own progress and activity. These factors could possibly also contribute to achieving a higher level of activity.

Both groups of patients walked about 5000 to 6000 steps a day at steady state in the fifth week after discharge. The American Heart Association recommends that healthy subjects walk 10,000 steps a day for overall better health outcomes, including cardiovascular outcome [39]. The official recommendation by the American Heart Association and World Health Organization is 150 minutes of moderate-intensity aerobic physical activity a day for 5 days a week [40], equivalent to 7000-8000 steps a day. Prior studies evaluating the number of steps in patients with coronary artery disease in secondary prevention proposed a target of 7500 steps a day to correlate with improved condition in terms of lipid profiles, muscle endurance, and body mass index [41,42].

Limitations

This explorative observational study has multiple limitations. The lack of validation for this wearable technology was already described in the Discussion section.

Subjects were scheduled for either OPCAB or RA-MIDCAB surgery based on the coronary anatomy (number of grafts), comorbidities, and endovascular options. Therefore, they were matched according to baseline characteristics, cardiac history, and comorbidity profile, but the groups were found to be significantly different for body mass index, which was higher in the OPCAB group. Although body mass index was not significantly correlated to the number of steps at any point of the study, it might still be an influencing factor for physical activity. However, the difference was not taken into account in further analysis.

As stated above, physical activity is influenced by other factors in addition to the impact of a surgical intervention. Not all patients are equally active in nature and the differences may depend on self-efficacy, the choice for physiotherapy, and cardiac rehabilitation thereafter. Physiotherapy in the first weeks after surgery and the following Phase II cardiac rehabilitation may significantly influence the progress in physical activity.

Subjects were recommended to wear the activity tracker all day and to take it off only for charging or while showering. To verify if patients were constantly wearing the device, we checked for any missing data in continuous heart rate monitoring. Apart from this measure, it was difficult to supervise the wearing time. In an ideal setting, these patient cohorts would be analyzed in a randomized controlled trial. In addition, the present study is based on a small sample size. Hence, the results should be interpreted with caution, and further investigations should be carried out before outlining definitive conclusions.

Conclusions

This research aimed to evaluate postsurgical cardiac rehabilitation progress by using commercially available wearable technology. We confirm our hypothesis that RA-MIDCAB patients have an advantage over OPCAB patients with regard to revalidation. Although not statistically significant, the RA-MIDCAB patient cohort showed a clear trend towards higher physical activity level in the first weeks after surgery. The exact hinge point must be confirmed with a larger number of patients. This work highlighted the feasibility of the use of wearable technology for physical activity monitoring in a clinical setting. Further research should be conducted to evaluate the accuracy and reliability of wearable technology before it serves clinical applications, especially in nonhealthy subjects with an altered gait pattern.

Conflicts of Interest

None of the authors have any conflict of interest to declare with regard to the design or execution of this study. The Fitbit trackers were purchased by the Department of Cardiac Surgery of the University Hospitals Leuven.

Multimedia Appendix 1

Inclusion and exclusion criteria.

[PDF File (Adobe PDF File), 15KB - [mhealth_v7i1e9865_app1.pdf](#)]

Multimedia Appendix 2

Baseline characteristics and demographics.

[PDF File (Adobe PDF File), 11KB - [mhealth_v7i1e9865_app2.pdf](#)]

Multimedia Appendix 3

Spearman correlations with the number of steps in weeks 1, 2, and 5.

[PDF File (Adobe PDF File), 13KB - [mhealth_v7i1e9865_app3.pdf](#)]

References

1. Cheng TO. History of coronary artery bypass surgery--half of a century of progress. *Int J Cardiol* 2012 May 17;157(1):1-2. [doi: [10.1016/j.ijcard.2012.02.018](https://doi.org/10.1016/j.ijcard.2012.02.018)] [Medline: [22425221](https://pubmed.ncbi.nlm.nih.gov/22425221/)]

2. Soylyu E, Harling L, Ashrafian H, Attaran S, Athanasiou C, Punjabi P, et al. A systematic review of the safety and efficacy of distal coronary artery anastomotic devices in MIDCAB and TECAB surgery. *Perfusion* 2016 Oct;31(7):537-543. [doi: [10.1177/0267659115618004](https://doi.org/10.1177/0267659115618004)] [Medline: [26590391](https://pubmed.ncbi.nlm.nih.gov/26590391/)]
3. Ruel M, Une D, Bonatti J, McGinn J. Minimally invasive coronary artery bypass grafting: is it time for the robot? *Curr Opin Cardiol* 2013 Nov;28(6):639-645. [doi: [10.1097/HCO.0b013e3283653fd1](https://doi.org/10.1097/HCO.0b013e3283653fd1)] [Medline: [24077608](https://pubmed.ncbi.nlm.nih.gov/24077608/)]
4. Rogers C, Pike K, Angelini G, Reeves B, Glauber M, Ferrarini M, et al. An open randomized controlled trial of median sternotomy versus anterolateral left thoracotomy on morbidity and health care resource use in patients having off-pump coronary artery bypass surgery: the Sternotomy Versus Thoracotomy (STET) trial. *J Thorac Cardiovasc Surg* 2013 Aug;146(2):306-16.e1 [FREE Full text] [doi: [10.1016/j.jtcvs.2012.04.020](https://doi.org/10.1016/j.jtcvs.2012.04.020)] [Medline: [22944093](https://pubmed.ncbi.nlm.nih.gov/22944093/)]
5. Magee M, Mack M. Robotics and coronary artery surgery. *Curr Opin Cardiol* 2002 Nov;17(6):602-607. [Medline: [12466701](https://pubmed.ncbi.nlm.nih.gov/12466701/)]
6. Poston RS, Tran R, Collins M, Reynolds M, Connerney I, Reicher B, et al. Comparison of economic and patient outcomes with minimally invasive versus traditional off-pump coronary artery bypass grafting techniques. *Ann Surg* 2008 Oct;248(4):638-646 [FREE Full text] [doi: [10.1097/SLA.0b013e31818a15b5](https://doi.org/10.1097/SLA.0b013e31818a15b5)] [Medline: [18936577](https://pubmed.ncbi.nlm.nih.gov/18936577/)]
7. Niebauer J. Is There a Role for Cardiac Rehabilitation After Coronary Artery Bypass Grafting? Treatment After Coronary Artery Bypass Surgery Remains Incomplete Without Rehabilitation. *Circulation* 2016 Jun 14;133(24):2529-2537. [doi: [10.1161/CIRCULATIONAHA.116.021348](https://doi.org/10.1161/CIRCULATIONAHA.116.021348)] [Medline: [27297345](https://pubmed.ncbi.nlm.nih.gov/27297345/)]
8. Mendes M. Is There a Role for Cardiac Rehabilitation After Coronary Artery Bypass Grafting? There is No Role for Cardiac Rehabilitation After Coronary Artery Bypass Grafting. *Circulation* 2016 Jun 14;133(24):2538-2543. [doi: [10.1161/CIRCULATIONAHA.115.017800](https://doi.org/10.1161/CIRCULATIONAHA.115.017800)] [Medline: [27297346](https://pubmed.ncbi.nlm.nih.gov/27297346/)]
9. European Association of Cardiovascular Prevention Rehabilitation Committee for Science Guidelines, EACPR, Corrà U, Piepoli MF, Carré F, Heuschmann P, Document Reviewers, et al. Secondary prevention through cardiac rehabilitation: physical activity counselling and exercise training: key components of the position paper from the Cardiac Rehabilitation Section of the European Association of Cardiovascular Prevention and Rehabilitation. *Eur Heart J* 2010 Aug;31(16):1967-1974. [doi: [10.1093/eurheartj/ehq236](https://doi.org/10.1093/eurheartj/ehq236)] [Medline: [20643803](https://pubmed.ncbi.nlm.nih.gov/20643803/)]
10. Munos B, Baker P, Bot B, Crouthamel M, de Vries G, Ferguson I, et al. Mobile health: the power of wearables, sensors, and apps to transform clinical trials. *Ann N Y Acad Sci* 2016 Dec;1375(1):3-18. [doi: [10.1111/nyas.13117](https://doi.org/10.1111/nyas.13117)] [Medline: [27384501](https://pubmed.ncbi.nlm.nih.gov/27384501/)]
11. Kas K. *Nooit Meer Ziek*. Antwerpen: Van Halewyck Nv; 2015.
12. Piwek L, Ellis D, Andrews S, Joinson A. The Rise of Consumer Health Wearables: Promises and Barriers. *PLoS Med* 2016 Feb;13(2):e1001953 [FREE Full text] [doi: [10.1371/journal.pmed.1001953](https://doi.org/10.1371/journal.pmed.1001953)] [Medline: [26836780](https://pubmed.ncbi.nlm.nih.gov/26836780/)]
13. Huang K, Liu W, He D, Huang B, Xiao D, Peng Y, et al. Telehealth interventions versus center-based cardiac rehabilitation of coronary artery disease: A systematic review and meta-analysis. *Eur J Prev Cardiol* 2015 Aug;22(8):959-971. [doi: [10.1177/2047487314561168](https://doi.org/10.1177/2047487314561168)] [Medline: [25488550](https://pubmed.ncbi.nlm.nih.gov/25488550/)]
14. Frederix I, Vanhees L, Dendale P, Goetschalckx K. A review of telerehabilitation for cardiac patients. *J Telemed Telecare* 2015 Jan;21(1):45-53. [doi: [10.1177/1357633X14562732](https://doi.org/10.1177/1357633X14562732)] [Medline: [25475219](https://pubmed.ncbi.nlm.nih.gov/25475219/)]
15. Kaiser D, Harrington R, Turakhia M. Wearable Fitness Trackers and Heart Disease. *JAMA Cardiol* 2016 Dec 01;1(2):239 [FREE Full text] [doi: [10.1001/jamacardio.2016.0354](https://doi.org/10.1001/jamacardio.2016.0354)] [Medline: [27437902](https://pubmed.ncbi.nlm.nih.gov/27437902/)]
16. Windecker S, Kolh P, Alfonso F, Collet J, Cremer J, Falk V, Grupa Robocza Europejskiego Towarzystwa Kardiologicznego (ESC), Europejskie Stowarzyszenie Chirurgii Serca i Klatki Piersiowej (EACTS) do spraw rewaskularyzacji mięśnia sercowego, European Association for Percutaneous Cardiovascular Interventions (EAPCI). [2014 ESC/EACTS Guidelines on myocardial revascularization]. *Kardiol Pol* 2014;72(12):1253-1379. [doi: [10.5603/KP.2014.0224](https://doi.org/10.5603/KP.2014.0224)] [Medline: [25524605](https://pubmed.ncbi.nlm.nih.gov/25524605/)]
17. Eapen Z, Peterson E. Can Mobile Health Applications Facilitate Meaningful Behavior Change?: Time for Answers. *JAMA* 2015;314(12):1236-1237. [doi: [10.1001/jama.2015.11067](https://doi.org/10.1001/jama.2015.11067)] [Medline: [26393844](https://pubmed.ncbi.nlm.nih.gov/26393844/)]
18. Leth S, Hansen J, Nielsen O, Dinesen B. Evaluation of Commercial Self-Monitoring Devices for Clinical Purposes: Results from the Future Patient Trial, Phase I. *Sensors (Basel)* 2017 Jan 22;17(1) [FREE Full text] [doi: [10.3390/s17010211](https://doi.org/10.3390/s17010211)] [Medline: [28117736](https://pubmed.ncbi.nlm.nih.gov/28117736/)]
19. Fokkema T, Kooiman T, Krijnen W, VAN DER Schans CP, DE Groot M. Reliability and Validity of Ten Consumer Activity Trackers Depend on Walking Speed. *Med Sci Sports Exerc* 2017 Dec;49(4):793-800. [doi: [10.1249/MSS.0000000000001146](https://doi.org/10.1249/MSS.0000000000001146)] [Medline: [28319983](https://pubmed.ncbi.nlm.nih.gov/28319983/)]
20. Lauritzen J, Muñoz A, Luis Sevillano J, Civit A. The usefulness of activity trackers in elderly with reduced mobility: a case study. *Stud Health Technol Inform* 2013;192:759-762. [Medline: [23920659](https://pubmed.ncbi.nlm.nih.gov/23920659/)]
21. Wong C, Mentis H, Kuber R. The bit doesn't fit: Evaluation of a commercial activity-tracker at slower walking speeds. *Gait Posture* 2018 Dec;59:177-181. [doi: [10.1016/j.gaitpost.2017.10.010](https://doi.org/10.1016/j.gaitpost.2017.10.010)] [Medline: [29049964](https://pubmed.ncbi.nlm.nih.gov/29049964/)]
22. Kooiman T, Dontje M, Sprenger S, Krijnen W, van der Schans CP, de Groot M. Reliability and validity of ten consumer activity trackers. *BMC Sports Sci Med Rehabil* 2015;7:24 [FREE Full text] [doi: [10.1186/s13102-015-0018-5](https://doi.org/10.1186/s13102-015-0018-5)] [Medline: [26464801](https://pubmed.ncbi.nlm.nih.gov/26464801/)]
23. Bai Y, Welk GJ, Nam YH, Lee JA, Lee JM, Kim Y, et al. Comparison of Consumer and Research Monitors under Semistructured Settings. *Med Sci Sports Exerc* 2016 Jan;48(1):151-158. [doi: [10.1249/MSS.0000000000000727](https://doi.org/10.1249/MSS.0000000000000727)] [Medline: [26154336](https://pubmed.ncbi.nlm.nih.gov/26154336/)]

24. An H, Jones G, Kang S, Welk G, Lee JM. How valid are wearable physical activity trackers for measuring steps? *Eur J Sport Sci* 2017 Apr;17(3):360-368. [doi: [10.1080/17461391.2016.1255261](https://doi.org/10.1080/17461391.2016.1255261)] [Medline: [27912681](https://pubmed.ncbi.nlm.nih.gov/27912681/)]
25. Evenson K, Goto M, Furberg R. Systematic review of the validity and reliability of consumer-wearable activity trackers. *Int J Behav Nutr Phys Act* 2015 Dec 18;12:159 [FREE Full text] [doi: [10.1186/s12966-015-0314-1](https://doi.org/10.1186/s12966-015-0314-1)] [Medline: [26684758](https://pubmed.ncbi.nlm.nih.gov/26684758/)]
26. Diaz K, Krupka D, Chang M, Peacock J, Ma Y, Goldsmith J, et al. Fitbit: An accurate and reliable device for wireless physical activity tracking. *Int J Cardiol* 2015 Apr 15;185:138-140 [FREE Full text] [doi: [10.1016/j.ijcard.2015.03.038](https://doi.org/10.1016/j.ijcard.2015.03.038)] [Medline: [25795203](https://pubmed.ncbi.nlm.nih.gov/25795203/)]
27. Berra K, Rippe J, Manson J. Making Physical Activity Counseling a Priority in Clinical Practice: The Time for Action Is Now. *JAMA* 2015;314(24):2617-2618. [doi: [10.1001/jama.2015.16244](https://doi.org/10.1001/jama.2015.16244)] [Medline: [26662069](https://pubmed.ncbi.nlm.nih.gov/26662069/)]
28. Lin JS, O'Connor E, Evans CV, Senger CA, Rowland MG, Groom HC. Behavioral counseling to promote a healthy lifestyle in persons with cardiovascular risk factors: a systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med* 2014 Oct 21;161(8):568-578. [doi: [10.7326/M14-0130](https://doi.org/10.7326/M14-0130)] [Medline: [25155549](https://pubmed.ncbi.nlm.nih.gov/25155549/)]
29. Chiauzzi E, Rodarte C, DasMahapatra P. Patient-centered activity monitoring in the self-management of chronic health conditions. *BMC Med* 2015 Apr 09;13:77 [FREE Full text] [doi: [10.1186/s12916-015-0319-2](https://doi.org/10.1186/s12916-015-0319-2)] [Medline: [25889598](https://pubmed.ncbi.nlm.nih.gov/25889598/)]
30. Savage P, Ades P. Pedometer step counts predict cardiac risk factors at entry to cardiac rehabilitation. *J Cardiopulm Rehabil Prev* 2008;28(6):370-7; quiz 378. [doi: [10.1097/HCR.0b013e31818c3b6d](https://doi.org/10.1097/HCR.0b013e31818c3b6d)] [Medline: [19008690](https://pubmed.ncbi.nlm.nih.gov/19008690/)]
31. Frederix I, Solmi F, Piepoli M, Dendale P. Cardiac telerehabilitation: A novel cost-efficient care delivery strategy that can induce long-term health benefits. *Eur J Prev Cardiol* 2017 Dec;24(16):1708-1717. [doi: [10.1177/2047487317732274](https://doi.org/10.1177/2047487317732274)] [Medline: [28925749](https://pubmed.ncbi.nlm.nih.gov/28925749/)]
32. Gong W, Cai J, Wang Z, Chen A, Ye X, Li H, et al. Robot-assisted coronary artery bypass grafting improves short-term outcomes compared with minimally invasive direct coronary artery bypass grafting. *J Thorac Dis* 2016 Mar;8(3):459-468 [FREE Full text] [doi: [10.21037/jtd.2016.02.67](https://doi.org/10.21037/jtd.2016.02.67)] [Medline: [27076941](https://pubmed.ncbi.nlm.nih.gov/27076941/)]
33. Rabindranauth P, Burns J, Vessey T, Mathiason M, Kallies K, Paramesh V. Minimally invasive coronary artery bypass grafting is associated with improved clinical outcomes. *Innovations (Phila)* 2014;9(6):421-426. [doi: [10.1097/IMI.0000000000000110](https://doi.org/10.1097/IMI.0000000000000110)] [Medline: [25438111](https://pubmed.ncbi.nlm.nih.gov/25438111/)]
34. Lapiere H, Chan V, Sohmer B, Mesana T, Ruel M. Minimally invasive coronary artery bypass grafting via a small thoracotomy versus off-pump: a case-matched study. *Eur J Cardiothorac Surg* 2011 Oct;40(4):804-810. [doi: [10.1016/j.ejcts.2011.01.066](https://doi.org/10.1016/j.ejcts.2011.01.066)] [Medline: [21393011](https://pubmed.ncbi.nlm.nih.gov/21393011/)]
35. Halkos M, Vassiliades T, Myung R, Kilgo P, Thourani V, Cooper W, et al. Sternotomy versus nonsternotomy LIMA-LAD grafting for single-vessel disease. *Ann Thorac Surg* 2012 Nov;94(5):1469-1477. [doi: [10.1016/j.athoracsur.2012.05.049](https://doi.org/10.1016/j.athoracsur.2012.05.049)] [Medline: [22776082](https://pubmed.ncbi.nlm.nih.gov/22776082/)]
36. Birla R, Patel P, Aresu G, Asimakopoulos G. Minimally invasive direct coronary artery bypass versus off-pump coronary surgery through sternotomy. *Ann R Coll Surg Engl* 2013 Oct;95(7):481-485 [FREE Full text] [doi: [10.1308/003588413X13629960047119](https://doi.org/10.1308/003588413X13629960047119)] [Medline: [24112493](https://pubmed.ncbi.nlm.nih.gov/24112493/)]
37. Blanchard C, Arthur H, Gunn E. Self-efficacy and outcome expectations in cardiac rehabilitation: associations with women's physical activity. *Rehabil Psychol* 2015 Feb;60(1):59-66. [doi: [10.1037/rep0000024](https://doi.org/10.1037/rep0000024)] [Medline: [25602362](https://pubmed.ncbi.nlm.nih.gov/25602362/)]
38. Sweet S, Fortier M, Strachan S, Blanchard C, Boulay P. Testing a Longitudinal Integrated Self-Efficacy and Self-Determination Theory Model for Physical Activity Post-Cardiac Rehabilitation. *Health Psychol Res* 2014 Jan 13;2(1):1008 [FREE Full text] [doi: [10.4081/hpr.2014.1008](https://doi.org/10.4081/hpr.2014.1008)] [Medline: [26973926](https://pubmed.ncbi.nlm.nih.gov/26973926/)]
39. Nocon M, Hiemann T, Müller-Riemenschneider F, Thalau F, Roll S, Willich S. Association of physical activity with all-cause and cardiovascular mortality: a systematic review and meta-analysis. *Eur J Cardiovasc Prev Rehabil* 2008 Jun;15(3):239-246. [doi: [10.1097/HJR.0b013e3282f55e09](https://doi.org/10.1097/HJR.0b013e3282f55e09)] [Medline: [18525377](https://pubmed.ncbi.nlm.nih.gov/18525377/)]
40. Global recommendations on physical activity for health Internet. World Health Organization URL: http://www.who.int/dietphysicalactivity/factsheet_recommendations/en/ [accessed 2018-03-02] [WebCite Cache ID [6xcEIUr36](https://www.webcitation.org/6xcEIUr36)]
41. Bäck M, Cider A, Gillström J, Herlitz J. Physical activity in relation to cardiac risk markers in secondary prevention of coronary artery disease. *Int J Cardiol* 2013 Sep 20;168(1):478-483. [doi: [10.1016/j.ijcard.2012.09.117](https://doi.org/10.1016/j.ijcard.2012.09.117)] [Medline: [23041099](https://pubmed.ncbi.nlm.nih.gov/23041099/)]
42. Houle J, Valera B, Gaudet-Savard T, Auclair A, Poirier P. Daily steps threshold to improve cardiovascular disease risk factors during the year after an acute coronary syndrome. *J Cardiopulm Rehabil Prev* 2013;33(6):406-410. [doi: [10.1097/HCR.0000000000000021](https://doi.org/10.1097/HCR.0000000000000021)] [Medline: [24104407](https://pubmed.ncbi.nlm.nih.gov/24104407/)]

Abbreviations

CAB: coronary artery bypass

CABG: coronary artery bypass grafting

EuroSCORE II: European System for Cardiac Operative Risk Evaluation

NYHA: New York Heart Association functional classification of heart failure

OPCAB: off-pump coronary artery bypass

RA-MIDCAB: robotically assisted minimally invasive direct coronary artery bypass

STEMI: ST-segment elevated myocardial infarction

Edited by G Eysenbach; submitted 17.01.18; peer-reviewed by J Goris, F Vogt, J Pietilä; comments to author 15.02.18; revised version received 13.03.18; accepted 10.12.18; published 24.01.19

Please cite as:

Thijs I, Fresiello L, Oosterlinck W, Sinnaeve P, Rega F

Assessment of Physical Activity by Wearable Technology During Rehabilitation After Cardiac Surgery: Explorative Prospective Monocentric Observational Cohort Study

JMIR Mhealth Uhealth 2019;7(1):e9865

URL: <http://mhealth.jmir.org/2019/1/e9865/>

doi: [10.2196/mhealth.9865](https://doi.org/10.2196/mhealth.9865)

PMID:

©Isabeau Thijs, Libera Fresiello, Wouter Oosterlinck, Peter Sinnaeve, Filip Rega. Originally published in JMIR Mhealth and Uhealth (<http://mhealth.jmir.org>), 24.01.2019. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR mhealth and uhealth, is properly cited. The complete bibliographic information, a link to the original publication on <http://mhealth.jmir.org/>, as well as this copyright and license information must be included.