

Review

Efficacy, Effectiveness, and Quality of Resilience-Building Mobile Health Apps for Military, Veteran, and Public Safety Personnel Populations: Scoping Literature Review and App Evaluation

Melissa Voth^{1,2}, BEd; Shannon Chisholm², BSc; Hannah Sollid², BSc; Chelsea Jones^{1,3,4}, PhD; Lorraine Smith-MacDonald^{1,2}, PhD; Suzette Brémault-Phillips^{1,2}, PhD

¹Heroes in Mind, Advocacy and Research Consortium, Faculty of Rehabilitation Medicine, University of Alberta, Edmonton, AB, Canada

²Department of Occupational Therapy, Faculty of Rehabilitation, University of Alberta, Edmonton, AB, Canada

³Leiden University Medical Centre, Leiden University, Leiden, Netherlands

⁴Operational Stress Injury Clinic, Alberta Health Services, Edmonton, AB, Canada

Corresponding Author:

Chelsea Jones, PhD

Heroes in Mind, Advocacy and Research Consortium

Faculty of Rehabilitation Medicine

University of Alberta

1-94 Corbett Hall

8205 - 114 Street Edmonton

Edmonton, AB, T6G 2G4

Canada

Phone: 1 7804920404

Email: cweiman@ualberta.ca

Abstract

Background: Military members (MMs) and public safety personnel (PSP) are vulnerable to occupational stress injuries because of their job demands. When MMs and PSP transition out of these professions, they may continue to experience mental health challenges. The development and implementation of resilience-building mobile health (mHealth) apps as an emergent mental health intervention platform has allowed for targeted, cost-effective, and easily accessible treatment when in-person therapy may be limited or unavailable. However, current mHealth app development is not regulated, and often lacks both clear evidence-based research and the input of health care professionals.

Objective: This study aims to evaluate the evidence-based quality, efficacy, and effectiveness of resilience-building mobile apps targeted toward the MMs, PSP, and veteran populations via a scoping literature review of the current evidence base regarding resilience apps for these populations and an evaluation of free resilience apps designed for use among these populations.

Methods: The studies were selected using a comprehensive search of MEDLINE, CINAHL Plus, PsycINFO, SocINDEX, Academic Search Complete, Embase, and Google and were guided by PRISMA-ScR (Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews). A narrative synthesis of the resulting papers was performed. The Alberta Rating Index for Apps was used to conduct a review of each of the identified apps. The inclusion criteria consisted of apps that were free to download in either the Google Play Store or the Apple App Store; updated within the last 3 years; available in English and in Canada; and intended for use by MMs, veterans, and PSP.

Results: In total, 22 apps met the inclusion criteria for evaluation. The resilience strategies offered by most apps included psychoeducation, mindfulness, cognitive behavioral therapy, and acceptance and commitment therapy. Overall, 50% (11/22) of apps had been tested in randomized controlled trials, 7 (32%) apps had been evaluated using other research methods, and 5 (23%) apps had not been studied. Using the Alberta Rating Index for Apps, the app scores ranged from 37 to 56 out of 72, with higher rated apps demonstrating increased usability and security features.

Conclusions: The mHealth apps reviewed are well-suited to providing resilience strategies for MMs, PSP, and veterans. They offer easy accessibility to evidence-based tools while working to encourage the use of emotional and professional support with safety in mind. Although not intended to function as a substitute for professional services, research has demonstrated that mHealth apps have the potential to foster a significant reduction in symptom severity for posttraumatic stress disorder, depression, anxiety,

and other mental health conditions. In clinical practice, apps can be used to supplement treatment and provide clients with population-specific confidential tools to increase engagement in the treatment process.

(*JMIR Mhealth Uhealth* 2022;10(1):e26453) doi: [10.2196/26453](https://doi.org/10.2196/26453)

KEYWORDS

occupational stress injury; trauma; mHealth; resilience; mental health; military; veteran; public safety personnel; OSI; PTSD; mental health intervention; mobile phone

Introduction

Background

Globally, military members (MMs) and public safety personnel (PSP), for example, correctional workers, dispatchers, firefighters, paramedics, and police officers, experience increased exposure to trauma and stress in their daily activities, which can affect their mental health and well-being [1,2]. PSP and MMs are at an increased risk of developing occupational stress injuries (OSIs), including posttraumatic stress disorder (PTSD), major depressive disorder, generalized anxiety disorder, and increased anger, aggression, or hostility, which can lead to other challenges, such as substance abuse, relationship difficulties, and workplace absenteeism [1].

MMs in the Canadian Armed Forces (CAF) are at greater risk of mental health disorders and suicide risk compared with the Canadian civilian population [3]. Of the regular force CAF members, 32.2% self-reported a mental health problem related to emotions, stress, substances, or family in 2013-2014 [4]. In addition, within the 3-year period from 2013 to 2016, mental health conditions in the veteran populations showed an increase from 25.4% to 30.3%, with PTSD being the most commonly identified OSI [5]. A 2018 study indicated that across the Canadian PSP groups, 44% screened positive for at least one mental health disorder [1]. This study also found that 36.7% of surveyed Canadian police officers in particular screened positive for mental health conditions, primarily PTSD [1]. These populations face challenges related to attaining professional mental health, including displacement owing to relocation, working in remote geographic locations, and shift work.

Owing to the need for mental health support among MMs, veterans, and PSP, mobile health (mHealth) apps have emerged as a portable treatment modality option [6-8]. Interest and use of mHealth by clinicians has increased in recent years in health

care practice [9]. The latest estimates suggest that there are between 165,000 and 325,000 health and wellness apps currently available for download [10,11]. When considering the MMs, veteran, and PSP populations, mHealth apps have gained popularity as a mental health treatment modality because of their low costs, easy access, and in-the-moment interventions [12].

Resilience

Evidence illustrates that resilience training and interventions, primarily those focused on coping skills and self-efficacy, can work to support a decrease in psychological distress and symptoms of PTSD [13-16]. Resilience is a broad and often complex concept, which scholars have uniquely interpreted depending on the context and can encompass both the individual and the group [17]. For this study, we have defined resilience as follows: “The dynamic process of overcoming adverse experiences through the use of internal and external resources in order to foster healthy psychological functioning” [13,17-19].

Resilience has been identified as an important factor that enables individuals adapt to and recover from emotionally, physically, and psychologically distressing situations and trauma [17,20]. There have been a multitude of resilience models and frameworks proposed in recent years specific to military, veteran, and PSP populations. One such model was developed using a large meta-analysis study by the Defence Human Capability and Science Technology Centre in 2014 [21]. This was further refined by Precious and Lindsay [22] with the Australian Armed Forces, resulting in a pillar of the mental resilience model (Figure 1 [21,22]). This model collaboratively draws on the best evidence related to mental resilience, highlighting both aspects outside one’s locus of control (ie, learned skills, previous experience, and personality) and the activities and skills within one’s locus of control (ie, mental control, emotional regulation, coping, self-efficacy, sense of purpose, positive affect, and social support) [22].

Figure 1. Pillars of mental resilience [21,22].

The activities and skills listed in this model have been attributed to the fostering of resilience, including those used to improve emotional regulation and coping, such as mindfulness, grounding, and self-talk; positive affect, such as purposeful leisure activities; and interpersonal relationships [23,24]. According to Lopez [23], “resilient individuals have a greater likelihood of engaging in healthy and productive activities and having a better quality of life.” If an individual does not possess the self-regulatory abilities and tools necessary to deal with distressing situations, it can impact all their areas of life, including sleep, quality of life, work, motivation, interest, and engagement in daily activities [24]. Resilience is, therefore, vital for the MMs, veteran, and PSP populations, as it supports continued engagement in both purposeful activities and increases the ability to adapt to the challenges of daily living.

Digital Health

The term *digital health* refers to the use of electronic communication, services, and processes to deliver and facilitate health care services [25]. mHealth is a more recent subsegment of digital health that uses mobile technology to enable remote care and clinical health data collection. Digital health-based treatments have become a growing field of research and development for the MMs, PSP, and veteran populations. Popular modalities include virtual reality, web-based programs and games, and mHealth apps [26-28]. Current research indicates that these platforms are effective at reducing the symptoms of PTSD and other mental health disorders caused by exposure to trauma [28-32].

Specific to the MMs, PSP, and veteran populations, health care professionals (HCPs) are well-suited to use mHealth tools, such as apps, to supplement treatment and provide clients with immediate tools to help them overcome psychological impairment related to their traumas [9,31]. For example, if a military member or PSP encounters a stressful situation, the accessibility of apps can help them navigate their feelings in real time. In addition, mHealth apps may fill a gap for those who require mental health treatment but are faced with barriers,

such as stigma to visible help-seeking, long waiting times, a high mobility of their jobs, and geographic restrictions [8,30,33,34], all of which have been noted as problematic for the MMs, PSP, and veteran populations. Many apps are also cost-effective and may be beneficial where therapy services are limited or unavailable [35,36].

Although mHealth tools have significant potential, several barriers limit their full uptake in the health care system. For example, some forms of technology, though widely used, may still not be available to everyone under all circumstances, for example, locations with unreliable or reduced cellular service, limited Wi-Fi access, and financial barriers may impede their use [37]. More importantly, there is a paucity of peer-reviewed research published regarding mHealth apps to determine their uptake, impact, or the best practices for their development and regulation [12,30,37]. When considering the use and development of mHealth apps, the limited number of research studies and agencies available to regulate this field impacts the ability to meet the current needs of this rapidly expanding industry [12,37]. As a result, a significant proportion of apps currently available for download have limited evidence of the effectiveness, efficacy, and safety of their use. This can make it difficult for HCPs to identify the best mHealth resources to recommend to MMs, PSP, and veterans in support of their care.

Objectives

The aim of this study is to evaluate the evidence-based quality, efficacy, and effectiveness of resilience-building mobile apps targeted toward the MMs, PSP, and veteran populations. This is addressed through two objectives: (1) completion of a scoping literature review of the current evidence base regarding mental health apps for these populations and (2) evaluation of common free mental health apps designed for use among these populations. We then compare and triangulate the data from these 2 approaches. The determination of these factors will aid in improving the evidence base for mHealth apps to highlight their potential use for HCPs who may be providing mental health services.

Methods

Objective 1

A Scoping Literature Review

A scoping literature review was completed to explore the available literature on mHealth apps and their cultivation of resilience in MMs, PSP, and veteran populations. The PRISMA-ScR (Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews) was used to guide this scoping review and app search, both of which were conducted between December 18, 2020, and December 20, 2020 [38]. We selected the following electronic databases for the search: MEDLINE (Ovid interface), CINAHL Plus with Full Text (EBSCOhost interface), PsycINFO (Ovid interface), SocINDEX with Full Text (EBSCOhost interface), Academic Search Complete (EBSCOhost interface), and Embase (Ovid interface). We also used Google as a search tool to investigate gray literature in case it was not detected in the chosen database. Additional resources were manually selected to ensure a comprehensive retrieval of relevant studies that may

Textbox 1. Eligibility criteria for scoping literature review.

Inclusion criteria

- The search was limited to studies published from the year 2000 onward to include more current technology.
- Included articles focused on participants aged ≥ 16 years.
- Articles addressing resiliency, hardiness, or coping.

Exclusion criteria

- Data not pertaining to military populations or public service personnel.
- Studies published in languages other than English.
- No outcome of interest.

Definition of Search Terms

Keywords for the search were determined using three main concepts: specific population, resilience, and games (refer to [Multimedia Appendix 1](#) for a full description of the search terms).

Title and Abstract Screening

After the removal of duplicate articles from the search results, a minimum of 2 researchers screened each article based on their titles and abstracts to determine further eligibility for the literature review. Articles that did not meet the eligibility criteria were excluded. Conflicts were discussed and resolved via team consensus.

Full-Text Reads

The screened articles were then read in full by a minimum of 2 researchers. Conflicts were discussed and resolved via team meetings and final eliminations were made. The remaining articles were included in the scoping review.

have fallen outside of the predetermined search terms. The following steps were adhered to: (1) determination of the population, intervention, comparison, and outcomes research question, (2) determination of an eligibility criteria, (3) definition of search terms ([Multimedia Appendix 1](#)), (4) title and abstract screening, (5) full-text reading, (6) charting of the data, and (7) narrative synthesis. The final literature search took place on December 20, 2020.

Determination of Research Question

This literature review aimed to answer the following research question: What is the efficacy, effectiveness, and quality of mHealth apps on increasing resilience and self-regulatory strategies among MMs, PSP, and veterans?

Determination of Eligibility Criteria

The literature included in the search encompassed studies published from the year 2000 onward to account for the development of technology during this time. The articles had to address resilience or self-regulatory strategies. In addition, the articles were required to pertain to military, veteran, or PSP populations ([Textbox 1](#)).

Charting of the Data

The type of evidence, population, funding, interventions, outcomes, and recommendations were extracted from each remaining article and recorded on a spreadsheet.

Narrative Synthesis

A narrative synthesis was performed by 3 researchers to summarize the findings of the different studies and evaluation results. Narrative synthesis refers to an approach that relies primarily on the use of words and text to summarize and explain the findings of multiple studies associated with reviews [39]. Narrative synthesis can be particularly helpful when studies are heterogeneous and organizing the data in a more numerical or statistical format would be inappropriate. To conduct the narrative synthesis, 3 researchers first reviewed the included study results and deductively organized them using the pillars of mental resilience as a guide. Additional information related to study methods, key constructs, and study outcomes was then synthesized together to form a coherent understanding of each topic. Finally, the researchers provided a summary of the included articles and their relevance to app evaluations.

Objective 2: Evaluation of Available Mental Health Apps

The identified apps were chosen through the following steps: (1) identification of the apps addressed in the literature review, (2) establishment of eligibility criteria, and (3) search of the eligible apps by name in the Apple App Store or Google Play Store. Apps were then evaluated for overall quality using the Alberta Rating Index for Apps (ARIA) [40].

Textbox 2. Eligibility criteria for mobile health apps included in the study.

Inclusion criteria
<ul style="list-style-type: none"> • Apps that were available on the Apple App Store and Google Play Store. • Apps that were free to download. • Apps that were intended for use by military members or public safety personnel.
Exclusion criteria
<ul style="list-style-type: none"> • Apps that were not free to download. • Apps that were not available in the English language. • Apps that were not available in Canada. • Apps that were not yet released for public use or access.

Outcome Measure: ARIA

The ARIA ([Multimedia Appendix 2](#) [40]) was used as a measuring tool to rate each app included in the study [40]. There are two versions of the ARIA: one for care providers and one for end users. Although the ARIA has yet to be vigorously studied, its uptake by health care systems at the regional and national level has been swift, likely owing to its ease of use, applicability, and both clinical utility and accessibility to the client population. The ARIA is available in both English and French, allowing it to be used across Canada, particularly with all PSP and members of the CAF and Veterans Affairs Canada (VAC). The tool has 2 sections, A and B, with multiple feature items to be rated between 0 and 4: 0 being *strongly disagree* and 4 being *strongly agree*. Section A was completed before downloading the app. Items in section A included purpose, trustworthiness, privacy, and affordability [42]. Once completed, scores were added up for section A to obtain a total out of 24 [42]. Section B is scored once the user has spent a minimum of 10 minutes using the app [42]. The items in section B include security, trustworthiness, ease of use, functionality, target users, usefulness, and satisfaction [42]. When completed, the rater added up the section to obtain a score out of 48. Sections A and B were then added together to determine an overall score out of 72, with higher scores indicating better performance and user experience. The care-provider version also features a 0-4 rating scale on whether the app would be recommended to possible users. This measure does not impact the overall score of the app [42].

The care-provider version of the ARIA was chosen to review the selected apps because of its ability to be implemented by

Eligibility Criteria

Apps included in the study were available for download in the Apple App Store or Google Play Store, could be set up in English, and were accessible within the geographic region of Canada. Apps chosen were intended for use primarily with MMs, PSP, and veterans; however, app use and availability could also extend to civilian populations. We included only free apps because evidence indicates that although 93% of smartphone users are likely to download an app, only 35.8% would be inclined to pay for an app [41]. Refer to [Textbox 2](#) for the detailed eligibility criteria.

users, caregivers, and HCPs. Each identified app that met the eligibility criteria was evaluated by 2 reviewers using the ARIA. Conflicts were resolved through discussion with all researchers to determine the final ARIA score.


Results

Objective 1: Scoping Literature Review

Study Selection

After searching the databases and identifying records through additional sources, a total of 691 articles were identified. Seven additional records were identified from the other sources. After the removal of 252 duplicates, 63.5% (439/691) of articles remained for title and abstract screening. A total of 52.1% (360/691) of articles were determined to be irrelevant. Full-text reads were completed on the remaining 10.7% (74/691) of articles and 6.1% (42/691) of additional articles were excluded because of differences in outcomes of interest, irrelevancy to mHealth app use, limited qualitative or quantitative data, and repetitive publications. A total of 9 studies were excluded because they did not have a relevant outcome of interest. The reasons for exclusion were that the studies involved virtual reality without an app and were specifically for outcomes related to PTSD. From these 32 articles, 22 apps were identified as meeting the inclusion criteria. [Figure 2](#) shows a PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) summary chart of the study selection process. The results of the scoping review, including narrative synthesis, are described in subsequent sections.

Figure 2. A PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) chart for the systematic review study identification, selection, exclusion, and inclusion.



Study Findings

Evidence-Based Merit

Of the 22 apps identified in the scoping literature review, 11 (50%) apps had been tested in randomized controlled trials (RCTs), 7 (32%) apps using other methods, and 5 (23%) apps had not been research trialed at the time of this study. Alternative methods used to evaluate apps included a nonrandomized quasi-experimental pre-post follow-up design, pre- and posttest questionnaires, qualitative focus groups, and the Mobile App Rating Scale. Apps that were not research trialed were identified through scoping reviews. Two apps had been tested using multiple methods.

The literature indicated that out of the 22 apps, 15 (68%) used evidence-based strategies or incorporated evidence-based components within them. Among these 15 apps, 11 (73%) apps were developed using evidence-based practices as their foundation (*Virtual Hope Box*, *eQuoo*, *Mindfulness Coach*, *Resilience@Work/Mindarma*, *PE Coach*, *R2MR*, *High Res*, *PTSD Coach*, *CBT-I Coach*, *PHIT for Duty*, and *Stay Quit Coach*). In the RCT by Roy et al [28], 7 apps were included in

determining the effectiveness of symptom reduction (*Positive Activity Jackpot*, *Tactical Breather*, *Daily Yoga*, *Simply Yoga*, *Life Armor*, *PE Coach*, and *Virtual Hope Box*). However, it should be noted that apps were not separated but rather assessed as a group, rendering it difficult to identify the efficacy and effectiveness of each app individually. *Virtual Hope Box*, *eQuoo*, and *Resilience@Work/Mindarma* were evaluated separately in their respective RCT studies.

Mental Control, Emotional Regulation, Coping, and Self-efficacy

The resilience strategies offered by most of the apps fit within the pillars of mental resilience, including mindfulness training, psychoeducation, cognitive behavioral therapy, and acceptance and commitment therapy. Other strategies included biofeedback, sleep strategies, social engagement, mood tracking, time scheduling, and muscle relaxation techniques, such as yoga. Many of the apps included more than one strategy. Of the total 22 apps, 8 (36%) apps used mindfulness strategies, including *Resilience@Work/Mindarma*, *Mindfulness Coach*, *Daily Yoga*, *Simply Yoga*, *Virtual Hope Box*, *Tactical Breather*, *Breathe2Relax*, and *PHIT for Duty* and 4 (18%) apps

encompassed psychoeducation: *eQuoo*, *Resilience@Work/Mindarma*, *Life Armor*, and *PE Coach*. Apps that applied cognitive behavioral therapy techniques included *R2MR*, *High Res*, *PTSD Coach*, *Family Coach*, *CBT-I Coach*, *Stay Quit Coach*, and *eQuoo*. *Resilience@Work/Mindarma* was the only app in this study that drew from acceptance and commitment therapy principles.

Effect of Apps on Resilience

The evidence-based literature demonstrated that many of the apps increased resilience strategies for users as well as improved the overall aspects of mental health [28,30,43,44]. The *eQuoo* app was demonstrated to significantly improve the traits of resilience, personal growth, and positive relationships [40]. Similarly, *Resilience@Work/Mindarma* showed improved adaptive resilience and psychological flexibility [15]. Joyce et al [15] also found that this app significantly increased optimism and mindfulness practice among study participants. This study also found that the app increased use of emotional support from others and help-seeking behavior, which addresses the social support pillar of the pillars of mental resilience. In addition, *R2MR*, which stands for road to mental readiness, was found to be effective in increasing resilience and help-seeking behaviors in the participants and in reducing mental health stigma for individuals and entire workplaces [45]. Although much of the literature noted the integration of social support as a positive influence on app use, resilience and coping, and health promotion, there was an observed lack of social connectedness components within the apps reviewed [14,23,36,46].

Health Care and Social Support

Some apps, such as *PTSD Coach*, were demonstrated to be more effective in managing and reducing PTSD symptoms when used with the support of an HCP as opposed to independent use [43,47]. Sessions that provide instruction around optimal app use patterns as well as the app's purpose, can increase the user's knowledge and therefore adherence to treatment programs that

uses mHealth components [43,48,49]. Additional mentoring or coaching may also contribute to a greater elaboration on techniques introduced within apps and may enable the transfer of skills from the app experience [14]. Evidence further indicates that using mHealth apps with support as supplementary resources, rather than primary treatment, may enhance therapeutic outcomes and allow users more autonomy in their ability to track symptoms while sharing results with their providers [14,30,43]. When users complete treatment sessions, they have the ability to retain these tools for future use or reference to their care [30,50].

End User Preferences, Incentives, and Real-world Apps

Within the studies, there were important themes that arose through narrative synthesis around user preferences. One of those themes included apps that had a sense of progression, rhythm and routine, and elements of personal causation [43]. Having set challenges and a clear visualization of the progress helped increase app use, adherence to the intervention, and goal attainment, especially when users were able to establish their targeted goals beforehand [26,51]. Earning rewards increased the attractiveness of the app, as did receiving guidance and instant feedback on target behaviors [26]. Apps designed around games and narratives were often preferred as they encompassed many of these traits and were user-friendly and enjoyable [51]. However, many users also noted a preference for more practical application opportunities of target skills, so their learned behaviors could be transferred to real-world concerns [52].

Objective 2: Evaluation of Mental Health Apps

The ARIA scores ranged from 37 to 56 out of 72 (Multimedia Appendix 3). The highest overall scoring apps were *R2MR*, *PTSD Coach*, and *AIMS for Anger Management*, all of which had a total score of 56 (Figure 3). The lowest scoring apps were *Resilience@Work/Mindarma* (37 out of 72), *Breathe2Relax* (38 out of 72), and *High Res* (39 out of 72).

Figure 3. Highest scoring apps on Alberta Rating Index for Apps.



R2MR



PTSD Coach



AIMS for Anger Management

Certain items of the ARIA were identified as being particularly relevant to the MMs and PSP populations. These included security and confidentiality, trustworthiness, and usefulness and satisfaction. As security and confidentiality are a high priority for the MMs and PSP populations, the security item of the ARIA is imperative to acknowledge [30]. The following apps rated the highest on the security items with a 4 out of 4 rating on both

security and consent: *Virtual Hope Box*, *Positive Activity Jackpot*, *Daily Yoga*, and *Life Armor*. The apps that were rated 1 or less were *eQuoo*, *Resilience@Work/Mindarma*, *High Res*, and *PHIT for Duty*. Many apps were missing features such as a password or biometric identifier and instead relied on the user's phone security setup to provide these privacy measures.

Apps were rated higher on trustworthiness if developed by reliable sources with proof of evidence. In total, 15 apps were developed by government agencies in Canada and the United States, including the Department of Veteran Affairs, the Department of Defence, the National Centre for Telehealth and Technology, and VAC. For trustworthiness, *PTSD Family Coach*, *R2MR*, *Stay Quit Coach*, *CBT-i Coach*, and *eQuoo*, all scored 4 out of 4. Many apps did not state risk warnings associated with app use directly on their download page; however, information could occasionally be found within the app itself. For these cases, the apps were rated 1 out of 4 in trustworthiness. In the usefulness and satisfaction items, apps that rated the highest included *Mindfulness Coach* and *PTSD Coach*, both with combined scores of 11 out of 16 for the sections.

Discussion

Summary of Evidence

The aim of this study is to evaluate the evidence-based quality, efficacy, and effectiveness of resilience-building mobile apps targeted toward the MMs, PSP, and veteran populations. This involved the completion of a scoping literature review of the current evidence base regarding mental health apps for these populations and the evaluation of common free mental health apps designed for use among these populations. This study aims to provide some insight into the following research question: What is the efficacy, effectiveness, and quality of mHealth apps on increasing resilience and self-regulatory strategies among MMs, PSP, and veterans?

Overall, the results of this study indicated that most of the mHealth apps reviewed were well-suited to provide resilience strategies and skills for MMs, PSP, and veterans. These apps provided skills, strategies, and services, which could be categorized into the pillars of mental resilience and other commonly accepted definitions regarding psychoeducational interventions that can foster resilience. Common resilience strategies were well represented in many of the apps, often including mindfulness, psychoeducation, and positive coping or thinking skills. Our results indicated that no app fully addressed the 6 pillars of resilience identified by the armed forces of the United Kingdom and Australia. In total, 5 apps addressed 5 of the 6 pillars, whereas 12 apps addressed 4 or more of the 6 pillars, and 20 apps addressed 2 or more of the 6 pillars. The apps that rated the highest on the ARIA were *R2MR*, *Virtual Hope Box*, *eQuoo*, *Mindfulness Coach*, and *PTSD Family Coach*. The pillar most likely to be missing was *socialsupport*, with the apps largely ignoring this concept.

In the evaluation of the 22 apps, *R2MR*, *PTSD Coach*, and *AIMS for Anger Management* had the highest overall scores. Points were generally lost because of a missing statement of the risk of use of the app or a lack of security measures to protect app access from the individual's phone. Of the 22 apps assessed, 15 (68%) apps were developed by credible, military-focused government agencies (eg, VAC and the US Department of Veterans Affairs), which may help ensure that the content delivered was well-adapted for these populations. All but 2 of these apps were developed outside of Canada, which may impact

the accessibility to local resources and services owing to the geographically based content (eg, helplines). Future comparisons of the ARIA with other app evaluation tools, such as the Mobile App Rating Scale, may allow for a more in-depth understanding of mHealth apps; however, for the MMs and PSP populations, the ARIA's additional security and privacy questions provide a clearer understanding of population-specific concerns [7,44].

When considering evidence around the apps, it was noted that out of the 22 apps, only 11 (50%) apps had undergone evidence-based evaluation through an RCT. Although from the total 22 apps, only 11 (50%) apps were determined to be evidence-based, 15 (68%) apps had used evidence-based strategies or components within them. Some of the apps selected for evaluation in this study have not been evaluated in the evidence-based literature. This is partially a result of the large creation and turnover of mHealth apps available for download as well as the currently limited regulations guiding their development [7,12,30]. With this being an understudied area of research, a lack of evidence influenced other potentially important client considerations. For example, parameters around effective dosages of apps, such as how long and how often a user was required to use the tool to see lasting effects, were not addressed. Instead, many of the apps identified in the studies relied on user feedback to conclude whether the application was clinically effective [50]. Another usability component considered through the ARIA was the presentation of information. Many of the apps relied on large blocks of text to present educational information and, as noted by O'Toole and Brown [30], this format can be overwhelming for users and cause disadvantages for those with alternative learning styles. This review illustrates the limitations of both the evidence and the potential quality of the apps being proposed to support resilience in MMs, PSP, and veterans.

Although these apps incorporate strategies and skills that may assist in facilitating resilience, it must be acknowledged that there are other factors that impact their ability to increase foster resilience, such as individual motivation, education on use of the app, access to social support, and the use of apps together with an HCP or independently. In addition, care should be taken regarding the specific designation of these apps as *resilience apps* (particularly in light of the MMs, PSP, and veteran populations). As a universal operational definition of resilience is lacking, a clear understanding of what elements constitute or are most important to resilience is also lacking. To illustrate, the Canadian studies of resilience in MMs have independently and not always cohesively explored the constructs of personality, positive affect, mastery, and social support [53]; neuroticism, military hardiness, and problem-solving coping [54]; and conscientiousness, emotional stability, and positive social interactions [55]. This inability to effectively define what is meant by resilience within a specific organization, such as the CAF, is problematic for both the research and intervention development. This problem is compounded when exploring how the construct is defined by the militaries of other countries (ie, Australia, United Kingdom, and United States vs Canada) and how it may enhance mental health [56]. Without research to definitively measure resilience before and after app use and without an effective definition of resilience—which impairs the

researcher's ability to quantify the concept—it is not possible to decisively conclude whether these apps increase resilience.

The Role of HCPs in mHealth and Resilience

Mental health can impact the daily functioning of an individual, and the concept of resilience is closely tied to mental health and well-being [23]. MMs and PSP are more likely to be exposed to traumatic experiences and are well-suited to resilience interventions [4,5]. HCPs can support individuals in navigating the environment for external resources and addressing the barriers in multiple domains that they may be experiencing.

Smartphones and technology are part of daily habits in the modern era, and HCPs can identify how to incorporate mHealth apps into health care settings. As MMs, PSP, and veterans face many unique challenges in terms of sudden environmental, lifestyle, and role changes, mHealth tools can present a more feasible option for access [24]. In a clinical context, HCPs can recommend apps to provide ongoing support outside of therapy services, create a tool for sharing health information, increase engagement in the treatment process, and sustain benefits gained once the provided services end [28]. For the mHealth apps themselves, the inclusion of both clinicians and users in their development can both ensure strategies meet the needs of their clients, thus encouraging wider acceptability and utility and helping clinicians better identify evidence-based features [30,57,58].

HCPs have a responsibility to advocate for best practices; this can be challenging with mHealth because of the high rate of app development and the inability for evidence-based practice to keep up [12,30,37]. Using tools such as the ARIA can help clinicians determine an app's usability and evidence base; however, HCPs must also ensure that the app is a good fit for clients in terms of their interests, values, abilities, and routines. HCPs can then use this information to collaborate with the client, customize apps that meet the client's needs and interests, and promote engagement with the apps. Creating client autonomy through app literacy will allow users to take more control in choosing treatment methods and encourage greater client-centered practice [59]. It is important to note, however, that many of these apps perform best when combined with the services of HCPs. This may include providing learning sessions before use to increase efficacy and optimal use patterns, weekly phone check-ins, or using the apps to enhance existing therapeutic interactions between clinicians and users [43,48-50]. In much of the literature, gamification and the integration of social components strengthened mHealth app use and engagement, resulting in more positive outcomes and an increased sense of peer support for the MMs and PSP populations [14,36,46]; however, this was lacking in practice.

Strengths and Limitations

This study has several strengths. Both the scoping literature review and app evaluation were conducted following a planned a priori procedure, with attention to ensuring quality control and minimizing bias. The detailed search strategy in the literature review was extensive, including 6 databases. The inclusion and exclusion criteria were determined before the study onset and adhered to throughout for both the literature

review and app evaluation. We also used appropriate calibration and at least two independent reviewers for all stages of the process.

There are certain limitations to this study, which should also be acknowledged. In the literature review, only studies written in English were included. The app selection criteria were limited to apps available for download in Canada, which excluded potentially beneficial apps available in other geographic locations. In addition, as the researchers only had access to iPhones, the apps were not tested on Android devices, which could have an impact on the usability criteria.

Although the authors identified the ARIA as an appropriate evaluation tool to use, it should be acknowledged that this is still relatively new and has not yet been extensively researched, used, or validated at this point. As such, there are currently no similar studies conducted with the ARIA, with which the present results could be compared. The results listed within this study only reflect app use from the perspective of clinicians, which could create bias. It will be important in future studies to invite users from the MMs, PSP, and veteran populations to complete the *user version* of the ARIA.

Future Research and Directions

Regarding the future of mHealth and resilience, there is much work to do in the areas of research, development, and policy. First, despite its use in health care contexts, future research is required to determine how the ARIA scores correlate to app adherence, acceptance, and adoption by users as well as to health outcomes. Further comparison of the ARIA with other app evaluation tools would be valuable in understanding the utility, criterion validity, and other concepts related to its ability to rate apps from the perspective of HCPs and clients as end users.

As previously mentioned, there is a paucity of evidence-based studies on the existing apps geared toward both resilience and the specific patient population of MMs, PSP, and veterans. Although this lack of research does not necessarily indicate that apps are of poor quality, it highlights the need for further research on health app development to ensure safety, effectiveness, and efficacy. It has been identified that traditional study design and research methods may be inappropriate for the study of mHealth as technology evolves much faster than traditional evidence-based research [8,60]. The lack of empirical research to demonstrate the effectiveness of apps on resilience may be related to the short time frame during which mHealth apps have emerged and the speed at which their availability changes [8]. Innovative and novel research methods that can address the demands of mHealth and the rapidly changing world of app development and use are needed to assist with the quality control of these tools used by both HCPs and patient populations.

Another area of study related to mHealth that is important and specific to military and PSP users is privacy, security, and confidentiality. Although there is a higher expectation of privacy for apps that involve health care information, military and PSP organizations may be subject to other restrictions on internet access that may impede the use of the app or demand higher security. Data sharing and privacy are considerations that require

attention from researchers, HCPs, and the general public when deciding on which app to use or if app use is appropriate at all [8]. Future systems of app evaluation and research would benefit from adding a component that considers data sharing, storage, and privacy in highly sensitive and secure patient populations (ie, military and PSP).

Future initiatives to assist HCPs and their clients in navigating the world of mHealth would be an asset to balance client autonomy through app literacy and would assure that apps have some level of evidence-based merit. As mHealth use is on the rise among many HCP and client populations, training to use apps to support service delivery is of utmost importance for health care organizations. The establishment of clear practice guidelines is important for both HCPs and clients, so that expectations about the usefulness and effectiveness of the app are appropriately managed. Currently, clients may have overly enthusiastic ideas about the effectiveness of these apps to develop their resilience and support their mental health, even when seeking professional mental health treatment might be necessary. Similarly, issues of risk and safety in mental health apps (including resilience) also need to be addressed.

The question of what constitutes a resilience app versus, for example, a wellness app, is also a murky territory that requires further navigation. Until a universally operationalized definition has been established for resilience, it is likely that confusion will remain regarding the codification of specific resilience skills and strategies. For example, family- and community-level factors were seldom addressed in our identified apps, despite being indicated as an important component of resilience [44,46]. This lack of inclusion is more surprising given the strong evidence that social support is a strong contributor to psychological health [23,24], overall well-being, and quality of life [24]. Similarly, empowering others to use their skills for stress reduction, coping, and building self-efficacy has been demonstrated to foster a significant reduction in the severity of the symptoms of PTSD, depression, and anxiety [12,13,28,58], which also alludes to the idea that the identified apps could be classified in terms of specific mental disorders. Until researchers, app developers, the various military and PSP

organizations, and HCPs can agree on the definitions of these concepts, determining which intervention targets and affects resilience will remain elusive.

Finally, the evaluation of resilience apps will remain challenging for all stakeholders and can affect the quality of the product unless all stakeholders were included in the consultation process. With the exception of the apps designed by the United States Department of Defense of Veterans Affairs, it was difficult to determine if the end user's perspective was incorporated into the development of the app. A collaborative approach to development, using both expert and user input, has been noted in recent studies as an effective approach to increasing the success of apps [57,58,60]. Ideally, in the future, mHealth development should engage the end users' input to assist with contextualization, which may increase the app acceptance, usability, and feasibility in a multitude of health care and possibly military or PSP settings.

Conclusions

Resilience is often targeted by HCPs through interventions that strengthen social support systems, foster greater self-concept, encourage optimism, promote the ability to reflect, and build emotional strength. Although not intended to function as a substitute for professional services and interventions, mHealth apps have the potential to foster resilience and support a significant reduction in symptom severity for OSIs, including PTSD, depression, and anxiety, in populations affected by OSIs, such as MMs, veterans, and PSP. Apps provide easy accessibility to evidence-based tools and encourage users to initiate help-seeking behaviors when stigma or uncertainty may impede the use of direct care. In clinical practice, HCPs can assist clients in identifying apps that support their habits and values and bolster participation and engagement in activities of daily living. As accessible, novel, and evidence-based interventions and resources for fostering resilience and addressing mental health become available, MMs, veterans, and PSP may be able to facilitate their healing, recovery, and growth, which would have a positive effect on their families, communities, organizations, and the public they serve.

Acknowledgments

The authors wish to thank Dr Peyman Azad Khaneghah for offering the use of the Alberta Rating Index for Apps and providing support for its implementation. This study would not have been possible without the support of the Heroes in Mind, Advocacy, and Consortium research laboratory.

The Recipient is providing this report of the project results on an "as is, where is" basis and makes no representations or warranties, either express or implied, as to any matter including, without limitation, whether the project results or any part or aspect of the same will be capable of statutory protection, the existence or non-existence of competing technology, the condition, quality or freedom from error of the project results or any part thereof, any merchantability, or its fitness for any particular purpose and all warranties and conditions expressed or implied, statutory or otherwise are hereby disclaimed. Neither the Recipient nor its officers, directors, employees, students or agents will be liable for any direct, consequential or other damage suffered by anyone resulting from the development or use of the project results or any inventions, technology or product produced in the course of or using the project results. The user of this report and/or any project results contained in the report uses the report and/or project results at the user's own risk.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Literature search strategy for scoping review.

[\[DOCX File, 14 KB-Multimedia Appendix 1\]](#)

Multimedia Appendix 2

Alberta Rating Index for Apps care-provider and user versions.

[\[PDF File \(Adobe PDF File\), 733 KB-Multimedia Appendix 2\]](#)

Multimedia Appendix 3

Alberta Rating Index for Apps score table.

[\[PDF File \(Adobe PDF File\), 42 KB-Multimedia Appendix 3\]](#)

References

1. Carleton RN, Afifi TO, Turner S, Taillieu T, Duranceau S, LeBouthillier DM, et al. Mental disorder symptoms among public safety personnel in Canada. *Can J Psychiatry* 2018 Jan;63(1):54-64 [[FREE Full text](#)] [doi: [10.1177/0706743717723825](https://doi.org/10.1177/0706743717723825)] [Medline: [28845686](https://pubmed.ncbi.nlm.nih.gov/28845686/)]
2. Ricciardelli R, Czarnuch S, Carleton RN, Gacek J, Shewmake J. Canadian public safety personnel and occupational stressors: how PSP interpret stressors on duty. *Int J Environ Res Public Health* 2020 Jul 01;17(13):4736 [[FREE Full text](#)] [doi: [10.3390/ijerph17134736](https://doi.org/10.3390/ijerph17134736)] [Medline: [32630259](https://pubmed.ncbi.nlm.nih.gov/32630259/)]
3. Rusu C, Zamorski MA, Boulos D, Garber BG. Prevalence comparison of past-year mental disorders and suicidal behaviours in the Canadian armed forces and the Canadian general population. *Can J Psychiatry* 2016 Apr;61(1 Suppl):46S-55S [[FREE Full text](#)] [doi: [10.1177/0706743716628856](https://doi.org/10.1177/0706743716628856)] [Medline: [27270741](https://pubmed.ncbi.nlm.nih.gov/27270741/)]
4. Theriault F, Gabler K, Naicker K. Health and lifestyle information survey of Canadian forces personnel 2013/2014. Regular Force Report. 2016. URL: <https://www.canada.ca/content/dam/dnd-mdn/documents/health/health-and-lifestyle-survey-2013-2014.pdf> [accessed 2021-12-23]
5. Thompson J, Van Til L, Poirier A, Sweet J, McKinnon K, Pedler D. Health and well-being of Canadian armed forces veterans: findings from the 2013 life after service survey. Research Directorate, Veterans Affairs Canada. 2014. URL: <https://www.veterans.gc.ca/pdf/about-us/research-directorate/2013-survey-caf-health.pdf> [accessed 2021-12-23]
6. Beshai S, Carleton N. Peer support and crisis-focused psychological intervention programs in Canadian first responders: blue paper. University of Regina Collaborative Centre for Justice and Safety. 2016. URL: <https://www.cipsrt-icrtsp.ca/en/publication/peer-support-and-crisis-focused-psychological-intervention-programs-in-canadian-first-responders-blue-paper> [accessed 2021-12-23]
7. Tam-Seto L, Wood VM, Linden B, Stuart H. Perceptions of an AI-supported mobile app for military health in the Canadian armed forces. *Military Behavioral Health* 2020 Nov 13;9(3):247-254. [doi: [10.1080/21635781.2020.1838364](https://doi.org/10.1080/21635781.2020.1838364)]
8. Jones C, O'Toole K, Jones K, Brémault-Phillips S. Quality of psychoeducational apps for military members with mild traumatic brain injury: an evaluation utilizing the mobile application rating scale. *JMIR Mhealth Uhealth* 2020 Aug 18;8(8):e19807 [[FREE Full text](#)] [doi: [10.2196/19807](https://doi.org/10.2196/19807)] [Medline: [32808937](https://pubmed.ncbi.nlm.nih.gov/32808937/)]
9. Buijink AW, Visser BJ, Marshall L. Medical apps for smartphones: lack of evidence undermines quality and safety. *Evid Based Med* 2013 Jun;18(3):90-92. [doi: [10.1136/eb-2012-100885](https://doi.org/10.1136/eb-2012-100885)] [Medline: [22923708](https://pubmed.ncbi.nlm.nih.gov/22923708/)]
10. Carlo AD, Hosseini Ghomi R, Renn BN, Areán PA. By the numbers: ratings and utilization of behavioral health mobile applications. *NPJ Digit Med* 2019;2:54 [[FREE Full text](#)] [doi: [10.1038/s41746-019-0129-6](https://doi.org/10.1038/s41746-019-0129-6)] [Medline: [31304400](https://pubmed.ncbi.nlm.nih.gov/31304400/)]
11. Lewis TL, Boissaud-Cooke MA, Aungst TD, Eysenbach G. Consensus on use of the term "App" versus "Application" for reporting of mHealth research. *J Med Internet Res* 2014 Jul 17;16(7):e174; discussion e174 [[FREE Full text](#)] [doi: [10.2196/jmir.3460](https://doi.org/10.2196/jmir.3460)] [Medline: [25033233](https://pubmed.ncbi.nlm.nih.gov/25033233/)]
12. Kuhn E, Owen JE. Advances in PTSD treatment delivery: the role of digital technology in PTSD treatment. *Curr Treat Options Psych* 2020 Mar 27;7(2):88-102. [doi: [10.1007/s40501-020-00207-x](https://doi.org/10.1007/s40501-020-00207-x)]
13. Combat-related trauma, posttraumatic stress disorder, and moral injury in combat medics: a comprehensive survey of the literature and critical analysis. ProQuest. URL: <https://www.proquest.com/openview/ae7c04d8fdea0b1e660499b5ff4f8589/1?pq-origsite=gscholar&cbl=18750&diss=y> [accessed 2021-12-23]
14. de Visser EJ, Dorfman A, Chartrand D, Lamon J, Freedy E, Weltman G. Building resilience with the Stress Resilience Training System: design validation and applications. *WOR* 2016 Jul 05;54(2):351-366. [doi: [10.3233/wor-162295](https://doi.org/10.3233/wor-162295)]
15. Joyce S, Shand F, Lal TJ, Mott B, Bryant RA, Harvey SB. Resilience@Work mindfulness program: results from a cluster randomized controlled trial with first responders. *J Med Internet Res* 2019 Feb 19;21(2):e12894 [[FREE Full text](#)] [doi: [10.2196/12894](https://doi.org/10.2196/12894)] [Medline: [30777846](https://pubmed.ncbi.nlm.nih.gov/30777846/)]
16. Pallavicini F, Argenton L, Toniazzi N, Aceti L, Mantovani F. Virtual reality applications for stress management training in the military. *Aerospace Med Human Performance* 2016 Dec 01;87(12):1021-1030. [doi: [10.3357/amhp.4596.2016](https://doi.org/10.3357/amhp.4596.2016)]

17. Southwick SM, Bonanno GA, Masten AS, Panter-Brick C, Yehuda R. Resilience definitions, theory, and challenges: interdisciplinary perspectives. *Eur J Psychotraumatol* 2014 Oct;5 [FREE Full text] [doi: [10.3402/ejpt.v5.25338](https://doi.org/10.3402/ejpt.v5.25338)] [Medline: [25317257](https://pubmed.ncbi.nlm.nih.gov/25317257/)]
18. Masten AS. Resilience from a developmental systems perspective. *World Psychiatry* 2019 Feb;18(1):101-102 [FREE Full text] [doi: [10.1002/wps.20591](https://doi.org/10.1002/wps.20591)] [Medline: [30600628](https://pubmed.ncbi.nlm.nih.gov/30600628/)]
19. Readiness, resilience, growth. Government of Canada. URL: <https://army.gc.ca/en/3-canadian-division/readiness-resilience-growth.page> [accessed 2021-12-23]
20. Green KT, Hayward LC, Williams AM, Dennis PA, Bryan BC, Taber KH, Mid-Atlantic Mental Illness Research, Education/Clinical Center Workgroup, et al. Examining the factor structure of the Connor-Davidson Resilience Scale (CD-RISC) in a post-9/11 U.S. military veteran sample. *Assessment* 2014 Aug 27;21(4):443-451 [FREE Full text] [doi: [10.1177/1073191114524014](https://doi.org/10.1177/1073191114524014)] [Medline: [24586090](https://pubmed.ncbi.nlm.nih.gov/24586090/)]
21. Examination of the evidence for preparing, sustaining and enhancing psychological and physical well-being - psychological resilience in the UK Armed Forces. Institute for Employment Studies. 2014. URL: <https://www.employment-studies.co.uk/project/examination-evidence-preparing-sustaining-and-enhancing-psychological-and-physical-well> [accessed 2021-12-23]
22. Precious D, Lindsay A. Mental resilience training. *J R Army Med Corps* 2019 Apr 22;165(2):106-108. [doi: [10.1136/jramc-2018-001047](https://doi.org/10.1136/jramc-2018-001047)] [Medline: [30580281](https://pubmed.ncbi.nlm.nih.gov/30580281/)]
23. Lopez A. Posttraumatic stress disorder and occupational performance: building resilience and fostering occupational adaptation. *Work* 2011;38(1):33-38. [doi: [10.3233/WOR-2011-1102](https://doi.org/10.3233/WOR-2011-1102)] [Medline: [21248418](https://pubmed.ncbi.nlm.nih.gov/21248418/)]
24. Edgelow MM, MacPherson MM, Arnaly F, Tam-Seto L, Cramm HA. Occupational therapy and posttraumatic stress disorder: a scoping review. *Can J Occup Ther* 2019 Apr;86(2):148-157. [doi: [10.1177/0008417419831438](https://doi.org/10.1177/0008417419831438)] [Medline: [31014080](https://pubmed.ncbi.nlm.nih.gov/31014080/)]
25. Benefits of digital health: what is digital health? Canada Health Infoway. URL: <https://www.infoway-inforoute.ca/en/what-we-do/benefits-of-digital-health/what-is-digital-health> [accessed 2021-06-23]
26. Pusey M, Wong KW, Rappa NA. Resilience interventions using interactive technology: a scoping review. *Interactive Learn Environ* 2020 Jun 11:1-16. [doi: [10.1080/10494820.2020.1772837](https://doi.org/10.1080/10494820.2020.1772837)]
27. Rizzo A, Buckwalter JG, Forbell E, Reist C, Difede J, Rothbaum BO, et al. Virtual reality applications to address the wounds of war. *Psychiatric Annals* 2013 Mar;43(3):123-138. [doi: [10.3928/00485713-20130306-08](https://doi.org/10.3928/00485713-20130306-08)]
28. Roy MJ, Costanzo ME, Highland KB, Olsen C, Clayborne D, Law W. An app a day keeps the doctor away: guided education and training via smartphones in subthreshold post traumatic stress disorder. *Cyberpsychol Behav Soc Netw* 2017 Aug;20(8):470-478. [doi: [10.1089/cyber.2017.0221](https://doi.org/10.1089/cyber.2017.0221)] [Medline: [28737954](https://pubmed.ncbi.nlm.nih.gov/28737954/)]
29. Bisson JI, van Deursen R, Hannigan B, Kitchiner N, Barawi K, Jones K, et al. Randomized controlled trial of multi-modular motion-assisted memory desensitization and reconsolidation (3MDR) for male military veterans with treatment-resistant post-traumatic stress disorder. *Acta Psychiatr Scand* 2020 Aug;142(2):141-151. [doi: [10.1111/acps.13200](https://doi.org/10.1111/acps.13200)] [Medline: [32495381](https://pubmed.ncbi.nlm.nih.gov/32495381/)]
30. O'Toole K, Brown CA. Evaluating the quality of resilience apps for military members and public safety personnel. *J Military Veteran Family Health* 2021 Feb 01;7(1):87-101. [doi: [10.3138/jmvfh-2020-0002](https://doi.org/10.3138/jmvfh-2020-0002)]
31. van Gelderen MJ, Nijdam M, Haagen J, Vermetten E. Interactive motion-assisted exposure therapy for veterans with treatment-resistant posttraumatic stress disorder: a randomized controlled trial. *Psychother Psychosom* 2020;89(4):215-227 [FREE Full text] [doi: [10.1159/000505977](https://doi.org/10.1159/000505977)] [Medline: [32203971](https://pubmed.ncbi.nlm.nih.gov/32203971/)]
32. mHealth App Developer Economics 2017: the current status and trends of the mHealth app market. Research2Guidance. URL: <https://research2guidance.com/> [accessed 2021-12-23]
33. Bush NE, Smolenski DJ, Denneson LM, Williams HB, Thomas EK, Dobscha SK. A virtual hope box: randomized controlled trial of a smartphone app for emotional regulation and coping with distress. *Psychiatr Serv* 2017 Apr 01;68(4):330-336. [doi: [10.1176/appi.ps.201600283](https://doi.org/10.1176/appi.ps.201600283)] [Medline: [27842473](https://pubmed.ncbi.nlm.nih.gov/27842473/)]
34. Vogt D. Mental health-related beliefs as a barrier to service use for military personnel and veterans: a review. *Psychiatr Serv* 2011 Feb;62(2):135-142. [doi: [10.1176/ps.62.2.pss6202_0135](https://doi.org/10.1176/ps.62.2.pss6202_0135)] [Medline: [21285091](https://pubmed.ncbi.nlm.nih.gov/21285091/)]
35. Iribarren SJ, Cato K, Falzon L, Stone PW. What is the economic evidence for mHealth? A systematic review of economic evaluations of mHealth solutions. *PLoS One* 2017;12(2):e0170581 [FREE Full text] [doi: [10.1371/journal.pone.0170581](https://doi.org/10.1371/journal.pone.0170581)] [Medline: [28152012](https://pubmed.ncbi.nlm.nih.gov/28152012/)]
36. Johnson D, Deterding S, Kuhn K, Staneva A, Stoyanov S, Hides L. Gamification for health and wellbeing: a systematic review of the literature. *Internet Interv* 2016 Nov;6:89-106 [FREE Full text] [doi: [10.1016/j.invent.2016.10.002](https://doi.org/10.1016/j.invent.2016.10.002)] [Medline: [30135818](https://pubmed.ncbi.nlm.nih.gov/30135818/)]
37. Imlawi J. Health website success: user engagement in health-related websites. *Int J Interact Mob Technol* 2017 Nov 27;11(6):49. [doi: [10.3991/ijim.v11i6.6959](https://doi.org/10.3991/ijim.v11i6.6959)]
38. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009 Jul 21;6(7):e1000097 [FREE Full text] [doi: [10.1371/journal.pmed.1000097](https://doi.org/10.1371/journal.pmed.1000097)] [Medline: [19621072](https://pubmed.ncbi.nlm.nih.gov/19621072/)]
39. Popay J, Roberts H, Sowden A, Petticrew M, Arai L, Rodgers M, et al. Guidance on the Conduct of Narrative Synthesis in Systematic Reviews: A Product From the ESRC Methods Programme. Lancaster: Lancaster University; 2006.

40. Azad-Khaneghah P, Roduta Roberts M, Stroulia E, Ferguson-Pell M, Liu L. Alberta Rating Index for Apps (ARIA): helping older adults find acceptable mobile health apps. *Gerontechnol* 2020 Oct 03;19:1-1. [doi: [10.4017/gt.2020.19.s.69884](https://doi.org/10.4017/gt.2020.19.s.69884)]
41. Kertz S, MacLaren Kelly J, Stevens K, Schrock M, Danitz S. A review of free iPhone applications designed to target anxiety and worry. *J Technol Behav Sci* 2017 Jan 11;2(2):61-70 [FREE Full text] [doi: [10.1007/s41347-016-0006-y](https://doi.org/10.1007/s41347-016-0006-y)]
42. Azad Khaneghah P. Alberta Rating Index for Apps (ARIA): an index to rate the quality of mobile health applications. University of Alberta Education & Research Archive. 2020. URL: <https://era.library.ualberta.ca/items/3c62ca2d-d7a1-4808-b1c0-4a60b05eeca> [accessed 2021-12-23]
43. Reyes AT, Serafica R, Sojobi A. College student veterans' experience with a mindfulness- and acceptance-based mobile app intervention for PTSD: a qualitative study. *Arch Psychiatr Nurs* 2020 Dec;34(6):497-506. [doi: [10.1016/j.apnu.2020.09.005](https://doi.org/10.1016/j.apnu.2020.09.005)] [Medline: [33280672](https://pubmed.ncbi.nlm.nih.gov/33280672/)]
44. Stoyanov SR, Hides L, Kavanagh DJ, Zelenko O, Tjondronegoro D, Mani M. Mobile app rating scale: a new tool for assessing the quality of health mobile apps. *JMIR Mhealth Uhealth* 2015 Mar 11;3(1):e27 [FREE Full text] [doi: [10.2196/mhealth.3422](https://doi.org/10.2196/mhealth.3422)] [Medline: [25760773](https://pubmed.ncbi.nlm.nih.gov/25760773/)]
45. Szeto A, Dobson KS, Knaak S. The road to mental readiness for first responders: a meta-analysis of program outcomes. *Can J Psychiatry* 2019 Jun;64(1_suppl):18S-29S [FREE Full text] [doi: [10.1177/0706743719842562](https://doi.org/10.1177/0706743719842562)] [Medline: [31010293](https://pubmed.ncbi.nlm.nih.gov/31010293/)]
46. Litvin S, Saunders R, Maier MA, Lüttke S. Gamification as an approach to improve resilience and reduce attrition in mobile mental health interventions: a randomized controlled trial. *PLoS One* 2020;15(9):e0237220 [FREE Full text] [doi: [10.1371/journal.pone.0237220](https://doi.org/10.1371/journal.pone.0237220)] [Medline: [32877425](https://pubmed.ncbi.nlm.nih.gov/32877425/)]
47. Possemato K, Kuhn E, Johnson E, Hoffman JE, Owen JE, Kanuri N, et al. Using PTSD Coach in primary care with and without clinician support: a pilot randomized controlled trial. *Gen Hosp Psychiatry* 2016;38:94-98. [doi: [10.1016/j.genhosppsy.2015.09.005](https://doi.org/10.1016/j.genhosppsy.2015.09.005)] [Medline: [26589765](https://pubmed.ncbi.nlm.nih.gov/26589765/)]
48. Roy M, Highland KB, Costanzo MA. GETSmart: Guided Education and Training via Smart phones to promote resilience. *Stud Health Technol Inform* 2015;219:123-128. [Medline: [26799892](https://pubmed.ncbi.nlm.nih.gov/26799892/)]
49. Weltman G, Lamon J, Freedy E, Chartrand D. Police department personnel stress resilience training: an institutional case study. *Glob Adv Health Med* 2014 Mar;3(2):72-79 [FREE Full text] [doi: [10.7453/gahmj.2014.015](https://doi.org/10.7453/gahmj.2014.015)] [Medline: [24808985](https://pubmed.ncbi.nlm.nih.gov/24808985/)]
50. Tam-Seto L, Wood VM, Linden B, Stuart H. A scoping review of mental health mobile apps for use by the military community. *Mhealth* 2018;4:57 [FREE Full text] [doi: [10.21037/mhealth.2018.12.01](https://doi.org/10.21037/mhealth.2018.12.01)] [Medline: [30701175](https://pubmed.ncbi.nlm.nih.gov/30701175/)]
51. Puskar K, Sun R, Gleeson A, Lampi T, Nichols D, Khan N. MilTeenChat app to promote coping resilience in military youth. *J Mil Veterans Health* Jan 2018;26(1):44-46. [doi: [10.3316/informit.553541068174068](https://doi.org/10.3316/informit.553541068174068)]
52. Thomas KH, Taylor SP, Hamner K, Glazer J, Kaufman E. Multi-site programming offered to promote resilience in military Veterans. *Calif J Health Promot* 2015 Sep 01;13(2):15-24. [doi: [10.32398/cjhp.v13i2.1820](https://doi.org/10.32398/cjhp.v13i2.1820)]
53. Lee JE, Sudom KA, McCreary DR. Higher-order model of resilience in the Canadian forces. *Canadian Journal of Behavioural Science / Revue canadienne des sciences du comportement* 2011 Jul;43(3):222-234. [doi: [10.1037/a0024473](https://doi.org/10.1037/a0024473)]
54. Skomorovsky A, Stevens S. Testing a resilience model among Canadian forces recruits. *Military Medicine* 2013 Aug;178(8):829-837. [doi: [10.7205/milmed-d-12-00389](https://doi.org/10.7205/milmed-d-12-00389)]
55. Lee JE, Sudom KA, Zamorski MA. Longitudinal analysis of psychological resilience and mental health in Canadian military personnel returning from overseas deployment. *J Occup Health Psychol* 2013 Jul;18(3):327-337. [doi: [10.1037/a0033059](https://doi.org/10.1037/a0033059)] [Medline: [23834447](https://pubmed.ncbi.nlm.nih.gov/23834447/)]
56. van der Meulen E, van der Velden PG, van Aert RC, van Veldhoven MJ. Longitudinal associations of psychological resilience with mental health and functioning among military personnel: a meta-analysis of prospective studies. *Soc Sci Med* 2020 Jun;255:112814. [doi: [10.1016/j.socscimed.2020.112814](https://doi.org/10.1016/j.socscimed.2020.112814)] [Medline: [32388075](https://pubmed.ncbi.nlm.nih.gov/32388075/)]
57. Deady M, Peters D, Lang H, Calvo R, Glozier N, Christensen H, et al. Designing smartphone mental health applications for emergency service workers. *Occup Med (Lond)* 2017 Aug 01;67(6):425-428. [doi: [10.1093/occmed/kqx056](https://doi.org/10.1093/occmed/kqx056)] [Medline: [28535246](https://pubmed.ncbi.nlm.nih.gov/28535246/)]
58. Colder Carras M, Kalbarczyk A, Wells K, Banks J, Kowert R, Gillespie C, et al. Connection, meaning, and distraction: a qualitative study of video game play and mental health recovery in veterans treated for mental and/or behavioral health problems. *Soc Sci Med* 2018 Nov;216:124-132 [FREE Full text] [doi: [10.1016/j.socscimed.2018.08.044](https://doi.org/10.1016/j.socscimed.2018.08.044)] [Medline: [30257787](https://pubmed.ncbi.nlm.nih.gov/30257787/)]
59. Peng W, Kanthawala S, Yuan S, Hussain SA. A qualitative study of user perceptions of mobile health apps. *BMC Public Health* 2016 Nov 14;16(1):1158 [FREE Full text] [doi: [10.1186/s12889-016-3808-0](https://doi.org/10.1186/s12889-016-3808-0)] [Medline: [27842533](https://pubmed.ncbi.nlm.nih.gov/27842533/)]
60. Mohr DC, Schueller SM, Riley WT, Brown CH, Cuijpers P, Duan N, et al. Trials of intervention principles: evaluation methods for evolving behavioral intervention technologies. *J Med Internet Res* 2015 Jul 08;17(7):e166 [FREE Full text] [doi: [10.2196/jmir.4391](https://doi.org/10.2196/jmir.4391)] [Medline: [26155878](https://pubmed.ncbi.nlm.nih.gov/26155878/)]

Abbreviations

- ARIA:** Alberta Rating Index for Apps
CAF: Canadian Armed Forces
HCP: health care professional

mHealth: mobile health

MM: military member

OSI: occupational stress injury

PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses

PRISMA-ScR: Preferred Reporting Items for Systematic Reviews and Meta-Analyses extension for Scoping Reviews

PSP: public safety personnel

PTSD: posttraumatic stress disorder

RCT: randomized controlled trial

VAC: Veterans Affairs Canada

Edited by L Buis; submitted 28.06.21; peer-reviewed by A Teles, J Austin; comments to author 01.09.21; revised version received 09.10.21; accepted 19.11.21; published 19.01.22

Please cite as:

Voth M, Chisholm S, Sollid H, Jones C, Smith-MacDonald L, Brémault-Phillips S

Efficacy, Effectiveness, and Quality of Resilience-Building Mobile Health Apps for Military, Veteran, and Public Safety Personnel Populations: Scoping Literature Review and App Evaluation

JMIR Mhealth Uhealth 2022;10(1):e26453

URL: <https://mhealth.jmir.org/2022/1/e26453>

doi: [10.2196/26453](https://doi.org/10.2196/26453)

PMID:

©Melissa Voth, Shannon Chisholm, Hannah Sollid, Chelsea Jones, Lorraine Smith-MacDonald, Suzette Brémault-Phillips. Originally published in JMIR mHealth and uHealth (<https://mhealth.jmir.org>), 19.01.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR mHealth and uHealth, is properly cited. The complete bibliographic information, a link to the original publication on <https://mhealth.jmir.org/>, as well as this copyright and license information must be included.