

Original Paper

Factors Associated With Self-reported Use of Web and Mobile Health Apps Among US Military Veterans: Cross-sectional Survey

Timothy P Hogan^{1,2,3}, PhD; Bella Etingen^{1,4}, PhD; Jessica M Lipschitz^{5,6}, PhD; Stephanie L Shimada^{1,2,7,8}, PhD; Nicholas McMahon^{1,2}, MPH; Derek Bolivar^{1,2}, BA; Felicia R Bixler^{1,4}, MS; Dawn Irvin^{1,4}, BS; Rachel Wacks^{1,2}, MPH, MA; Sarah Cutrona^{1,2,8}, MPH, MD; Kathleen L Frisbee^{1,9}, MPH, PhD; Bridget M Smith^{1,4,10}, PhD

¹eHealth Partnered Evaluation Initiative, Veterans Affairs Bedford Healthcare System, Bedford, MA, United States

²Center for Healthcare Organization and Implementation Research, Veterans Affairs Bedford Healthcare System, Bedford, MA, United States

³Peter O'Donnell Jr School of Public Health, University of Texas Southwestern Medical Center, Dallas, TX, United States

⁴Center of Innovation for Complex Chronic Healthcare, Hines Veterans Affairs Hospital, Hines, IL, United States

⁵Department of Psychiatry, Brigham and Women's Hospital, Boston, MA, United States

⁶Harvard Medical School, Boston, MA, United States

⁷Department of Health Law, Policy, and Management, Boston University School of Public Health, Boston, MA, United States

⁸Division of Health Informatics and Implementation Science, Department of Population and Quantitative Health Sciences, University of Massachusetts Chan Medical School, Worcester, MA, United States

⁹Office of Connected Care, Veterans Health Administration, US Department of Veterans Affairs, Washington, DC, United States

¹⁰Northwestern University Feinberg School of Medicine, Chicago, IL, United States

Corresponding Author:

Timothy P Hogan, PhD

Center for Healthcare Organization and Implementation Research

Veterans Affairs Bedford Healthcare System

200 Springs Road (152)

Building 70

Bedford, MA, 01730

United States

Phone: 1 781 687 3181

Email: timothy.hogan@va.gov

Abstract

Background: Despite their prevalence and reported patient interest in their use, uptake of health-related apps is limited. The Veterans Health Administration (VHA) has developed a variety of apps to support veterans; however, uptake remains low nationally.

Objective: We examined the prevalence of VHA health-related app use and how veterans learned about these apps in order to identify factors associated with their use.

Methods: As part of a VHA quality improvement initiative, we recruited a national cohort of veterans to obtain feedback on their use of technology for health and collected data from them via a cross-sectional survey. The survey data were supplemented with VHA administrative data. We used descriptive statistics to examine demographic and health characteristics, health-related technology use, and how veterans learned about apps. We assessed factors associated with app use using bivariate analyses and multiple logistic regression models.

Results: We had complete data on 1259 veterans. A majority of the sample was male (1069/1259, 84.9%), aged older than 65 years (740/1259, 58.8%), White (1086/1259, 86.3%), and non-Hispanic (1218/1259, 96.7%). Most respondents (1125/1259, 89.4%) reported being very comfortable and confident using computers, over half (675/1259, 53.6%) reported being an early adopter of technology, and almost half (595/1259, 47.3%) reported having used a VHA health-related app. Just over one-third (435/1259, 34.6%) reported that their VHA care team members encouraged them to use health-related apps. Respondents reported learning about available VHA health-related apps by reading about them on the VHA's patient portal (468/1259, 37.2%), being told about them by their VHA health care team (316/1259, 25.1%), and reading about them on the VHA's website (139/1259, 11%). Veterans who self-reported having used VHA health-related apps were more likely to receive care at the VHA (OR [odds ratio] 1.3, 95% CI 1.0-1.7), be in worse health (as assessed by Hierarchical Condition Community score; OR 1.1, 95% CI 1.0-1.2),

report owning a desktop or laptop computer (OR 1.8, 95% CI 1.1-3.1), have posttraumatic stress disorder (OR 1.4, 95% CI 1.1-1.9), and report having VHA health care team members encourage them to use the apps (OR 2.7, 95% CI 2.1-3.4).

Conclusions: We found strong associations between self-reported use by veterans of VHA health-related apps and multiple variables in our survey. The strongest association was observed between a veteran self-reporting app use and having received encouragement from their VHA health care team to use the apps. Veterans who reported receiving encouragement from their VHA care team members had nearly 3 times higher odds of using VHA apps than veterans who did not report receiving such encouragement. Our results add to growing evidence suggesting that endorsement of apps by a health care system or health care team can positively impact patient uptake and use.

(*JMIR Mhealth Uhealth* 2022;10(12):e41767) doi: [10.2196/41767](https://doi.org/10.2196/41767)

KEYWORDS

mobile health apps; patient engagement; consumer health informatics; provider encouragement; veterans

Introduction

There is an expanding number of apps available to help patients manage specific health conditions and promote overall well-being [1]. Evidence suggests use of health-related apps is associated with improved management of chronic health conditions [2] and mental health disorders [3,4], desirable health behavior change [5], and better medication adherence [6] and perceptions of health [7]. Studies have also shown that individuals are interested in using health-related apps to support self-management and to improve health [8]. Despite their prevalence and reported patient interest in using them, however, uptake of such apps remains limited [7,9,10].

A variety of barriers to health-related app adoption have been identified in previous work, including concerns about privacy and security [11-14], app usability issues [12,15], and limited proficiency with technology [14]. Recent literature also indicates that lack of awareness or knowledge of health-related apps is also a common barrier to adoption [14,16,17], highlighting the need to clarify how individuals who use health-related apps learn about them. Use of health-related apps has also been associated with certain patient demographic characteristics [18] and factors including positive perceptions of usefulness, motivation to change health behaviors or pursue a health goal, the availability of data visualization within the app, and the app not having any associated costs [9,11,14,16,18]. Importantly, research has also shown that the adoption of specific health-related apps may be bolstered if the app is recommended by a source that the target user trusts and finds credible [19].

US military veterans are an ideal population in which to examine the adoption of health-related apps. In comparison to the general US adult population, veterans often face significant health-related challenges, including disproportionate rates of physical and behavioral health diagnoses [20,21], and they commonly experience multiple comorbidities that require them to have frequent interaction with the health care system [22]. Some of these health concerns can be directly related to military service and difficulties with postservice community reintegration, while others represent common comorbidities experienced by US adults (eg, diabetes, heart disease, and chronic pain). For these reasons, US military veterans, like the broader population, stand to benefit from health-related apps and the support they offer for self-management and enhanced connection with health care providers and resources.

In recent years, studies have indicated that integrating apps into care for veterans may improve outcomes [23]. The effectiveness of multiple health-related apps targeted toward veterans has been demonstrated in randomized controlled trials, including use of the Virtual Hope Box app to support coping with negative emotions among veterans who experience suicidal ideation [24] and use of the PTSD Coach app to manage posttraumatic stress disorder (PTSD) symptom severity and increase PTSD treatment seeking [25]. As part of the Veterans Health Administration (VHA)'s digital health strategy, the VHA Office of Connected Care maintains a VHA app store, which contains a variety of mobile and web-based apps intended to support veterans in the management of their health. These apps are designed to address some of the unique needs of the veteran population, promote wellness and healthy behaviors, provide condition-specific self-management support, inform clinical management, and facilitate other transactions with the health care system that may be relevant to all veterans.

Despite the evolving evidence for their effectiveness, as well as recent literature indicating that many veterans are interested in using health-related apps [26,27] and have a device with which they can access them [28], uptake of VHA apps remains low nationally, and factors associated with veteran use of such apps are not well understood. Given that veterans represent a large patient population that could substantially benefit from the use of health-related apps, the objectives of this analysis were to examine the prevalence of VHA health-related app use, determine how veterans learned about these apps, and identify factors associated with their use.

Methods

Design

The VHA Office of Connected Care, in cooperation with investigators from the VHA Quality Enhancement Research Initiative program, developed the Veterans Engagement with Technology Collaborative (VET-C) cohort in 2017, the purpose of which was to engage veterans in the evaluation of VHA technologies that are intended to increase access, enhance coordination, and support self-management [28]. The VET-C cohort is a quality improvement resource that includes longitudinal survey data. Veterans who are part of the VET-C cohort are invited to provide feedback on their use of technologies for health, including VHA technologies, and this

feedback is used in turn to inform usability and broader uptake. We used the VET-C cohort to examine how veterans learned about VHA health-related apps and factors associated with their adoption.

Recruitment

To be eligible to join the VET-C cohort, veterans had to be users of VHA health care services, have a mobile phone, and be active users of the secure messaging feature of the VHA's online patient portal, My HealtheVet, a feature that is available to veterans who have a premium portal account. The secure messaging use requirement was intended as a proxy for receptivity to and use of VHA patient-facing technologies more generally. Active use of secure messaging was defined as having sent at least 5 secure messages to VHA clinical team members through the portal in the 12 months prior to cohort recruitment. Veterans who met these inclusion criteria were recruited from VHA facilities across the United States. VHA facilities were chosen as VET-C recruitment sites because they (1) had high rates of secure messaging, (2) served as field test sites for other new VHA patient-facing technologies, (3) were known to serve significant populations of women veterans and veterans from diverse ethnic and minority groups, and (4) had active research and evaluation programs.

Procedures

Recruitment lists to support development of the VET-C cohort were created by querying data from the VHA Corporate Data Warehouse (CDW). Veterans were called once by the evaluation team, and those who answered were told about the purpose of the VET-C cohort and invited to join. Veterans who were interested then completed the cohort baseline telephone survey, and evaluation team members entered their responses directly into an online REDCap database. During 2017 and 2018, 2727 veterans from 14 VHA facilities joined the VET-C cohort and completed the baseline survey. From March 2019 to March 2020, we administered a second survey to all veterans in the VET-C cohort who completed the baseline survey. This follow-up survey was completed by 1418 veterans in the cohort. Both our baseline and second surveys included validated question items, questions used in other studies, and new question items developed specifically for these surveys. They were developed in close consultation with leadership from the VHA Office of Connected Care, which is responsible for the health care system's digital health strategy. After we excluded veterans for whom there were missing data for the variables used in the analyses, we included a total of 1259 veterans in the current analyses.

Measures

For this paper, constructs of interest from the surveys included veteran demographics, health and health care use variables, technology ownership and use, VHA care team member encouragement to use VHA apps, and use of VHA health-related apps.

We asked all participants to respond to demographic questions on age, gender, race, ethnicity, relationship status, and education; report their perceived health status [29]; report where they normally received their medical care (the response options were

“mostly at the VHA,” “mostly outside the VHA,” “about half in the VHA, half outside the VHA,” and “nowhere”); and indicate the amount of time it typically took them to travel to their VHA primary care doctor from their home. We also asked participants about their technology ownership (ie, whether they owned a desktop computer, tablet computer, or mobile phone), whether they considered themselves to be early adopters of technology (ie, whether they liked to be among the first to get a new device, tech gadget, or app when it comes out), and how comfortable or confident they felt using computers (responses ranged from 0, “not at all,” to 5, “very”) [30].

Additionally, we asked participants to report whether they used VHA health-related apps and how they learned about the VHA health-related apps that are available. We also asked participants to report the perceived extent to which their VHA care team members encouraged them to use health-related apps (responses ranged from 1, “strongly disagree,” to 7, “strongly agree”).

We calculated the participants' prior-year comorbidity index as the Hierarchical Condition Community (HCC) score based on information in the VHA CDW [31,32]. The CDW was also used to identify diagnosed health conditions in the prior 5 years among our sample, and to fill in missing demographic data (eg, for age and gender).

Analyses

We used descriptive statistics (mean, range, and SD, or proportion, as appropriate) to characterize the demographics of the sample, as well as their reported health and health care use, technology ownership and use, VHA health-related app use, how they learned about VHA health-related apps, and their perceptions of VHA care team member encouragement to use health-related apps. We used bivariate analyses (the chi-square test and the *t* test) to examine differences among veterans who reported using (vs not using) VHA health-related apps. We then assessed factors associated with VHA health-related app use using unadjusted and adjusted multiple logistic regression models. We selected variables for inclusion in the unadjusted model based on significant bivariate associations with the outcome variables and known associations from the existing literature; we selected variables for inclusion in the adjusted model based on significant unadjusted associations at the $P < .1$ level and known associations from the existing literature. Statistical analyses were performed with Stata MP (version 14.2; StataCorp).

Ethics Approval

This work was reviewed by the institutional review boards of the VHA Bedford Healthcare System in Bedford, Massachusetts, and the Edward Hines Jr VHA Hospital in Hines, Illinois. The study was designated as a program evaluation for quality improvement purposes, exempting it from further oversight (VHA Handbook 1058.05).

Results

Sample Characteristics

Descriptive statistics and results of bivariate analyses are presented in Table 1. Overall, a majority of the sample was male

(1069/1259, 84.9%), aged over 65 years (740/1259, 58.8%), White (1086/1259, 86.3%), and non-Hispanic (1218/1259, 96.7%). Most (1125/1259, 89.4%) reported being very comfortable and confident using computers, and over half (675/1259, 53.6%) reported being an early adopter of technology. Almost half (595/1259, 47.3%) reported using VHA health-related apps. Just over one-third (435/1259, 34.6%) reported that their VHA care team members encouraged them to use health-related apps.

Respondents reported having learned about VHA apps through the VHA's patient portal (468/1259, 37.2%), their VHA health care team (316/1259, 25.1%), the VHA's government website (139/1259, 11%), veteran service organizations (102/1259, 8.1%), newsletters (66/1259, 5%), other veterans (64/1259, 5%), public app stores (64/1259, 5%), and the VHA mobile app store (58/1259, 5%). These results are presented in [Table 2](#).

Table 1. Comparison of demographic characteristics of veterans who were self-reported users or nonusers of health-related apps (N=1259).

Variable	Overall	Self-reported VHA ^a app users (595/1259; 47.3%)	Self-reported VHA app nonusers (664/1259; 52.7%)	P value
Demographics, n (%)				
Age older than 65 years	740 (58.8)	359 (60.3)	381 (57.4)	.29
Male	1069 (84.9)	505 (84.9)	564 (84.9)	.97
Race				.85
White	1086 (86.3)	512 (86.1)	574 (86.5)	
Black	118 (9.4)	55 (9)	63 (10)	
Other	55 (4)	28 (5)	27 (4)	
Hispanic ethnicity	41 (3)	24 (4)	17 (3)	.14
Relationship status: in a relationship ^b	906 (72)	426 (71.6)	480 (72.3)	.79
Education status: at least some college or vocational school ^c	1115 (88.6)	524 (88.1)	591 (89)	.60
Socioeconomic status ^d : “not very hard to pay for basics”	897 (71.3)	413 (69.4)	484 (72.9)	.17
Mostly receiving medical care at the VHA, n (%)	955 (75.9)	471 (79.2)	484 (72.9)	.009
Travel time to VHA ^e : more than 60 minutes, n (%)	198 (15.7)	82 (14)	116 (17.5)	.07
Perceived health status (fair/poor), n (%)	421 (33.4)	208 (35)	213 (32.1)	.28
Hierarchical Condition Community score, mean (SD)	1.6 (1.2)	1.7 (1.3)	1.5 (1.1)	.047
Technology use, n (%)				
Desktop or laptop computer	1184 (94)	571 (96)	613 (92.3)	.006
Tablet computer	721 (57.3)	356 (59.8)	365 (55)	.08
Smartphone	1123 (89.2)	541 (90.9)	582 (87.7)	.06
Early technology adopter	675 (53.6)	327 (55)	348 (52.4)	.37
Very comfortable or confident using computers	1125 (89.4)	540 (90.8)	585 (88.1)	.13
VHA health care team encouragement to use apps	435 (34.6)	276 (46.4)	159 (24)	<.001
Health conditions^f, n (%)				
Hypertension	726 (57.7)	345 (58)	381 (57.4)	.83
Osteoarthritis	716 (56.9)	347 (58.3)	369 (55.6)	.33
Diabetes	522 (41.5)	243 (40.8)	279 (42)	.67
Depression	515 (40.9)	252 (42.4)	263 (39.6)	.32
Chronic kidney disease	363 (28.8)	160 (26.9)	203 (30.6)	.15
Ischemic heart disease	340 (27)	165 (27.7)	175 (26.4)	.58
Asthma	302 (24)	140 (23.5)	162 (24.4)	.72
Posttraumatic stress disorder	269 (21.4)	148 (24.9)	121 (18.2)	.004
Peripheral vascular disease	202 (16)	103 (17.3)	99 (15)	.25
Anxiety disorders	201 (16)	102 (17.1)	99 (15)	.28
Atrial fibrillation	155 (12.3)	69 (12)	86 (13)	.47
Heart failure	118 (9.4)	62 (10)	56 (8)	.23
Acute myocardial infarction	115 (9.1)	48 (8)	67 (10)	.21
Stroke	90 (7)	45 (8)	45 (7)	.59
Prostate cancer	68 (5)	35 (6)	33 (5)	.48
Traumatic brain injury	63 (5)	33 (6)	30 (5)	.40
Chronic obstructive pulmonary disease	53 (4)	25 (4)	28 (4)	.99

Variable	Overall	Self-reported VHA ^a app users (595/1259; 47.3%)	Self-reported VHA app nonusers (664/1259; 52.7%)	P value
Colorectal cancer	20 (2)	9 (2)	11 (2)	.84
Lung cancer	17 (1)	7 (1)	10 (2)	.61

^aVHA: Veterans Health Administration.

^bDefined as married, in a civil union, or engaged; not being in a relationship was defined as being single, separated, divorced, or widowed.

^cDefined as 1 to 4 years of college or vocational school or a master's, professional, or doctoral degree.

^dDefined by the listed response to the question "How hard is it for you (and your family) to pay for the very basics like food and heating/cooling?"

^eResponse to the question "How many minutes does it usually take you to get to your healthcare practitioners office (your VHA primary care doctor's office)?"

^fIn the prior five years.

Table 2. Ways veterans reported having learned about Veterans Health Administration health-related apps. Respondents checked all options that applied to them.

Venue	Respondents (N=1259), n (%)
VHA ^a patient portal	468 (37.2)
VHA health care team members	316 (25.1)
VHA website	139 (11)
Veteran service organizations	102 (8.1)
Newsletters	66 (5)
Other veterans	64 (5)
Public app stores	64 (5)
VHA mobile app store	58 (5)
Other	17 (1)
Do not remember	12 (1)
At the hospital	12 (1)
VHA employee	7 (1)
Phone	6 (1)
Television	2 (0.2)

^aVHA: Veterans Health Administration.

Bivariate Comparisons of VHA Health-Related App Users and Nonusers

Bivariate analyses comparing respondents who reported using (vs not using) VHA health-related apps revealed that the former included greater proportions of veterans with PTSD (148/595, 24.9% vs 121/664, 18.2%; $P=.004$) and veterans who reported owning a desktop or laptop computer (571/595, 96% vs 613/664, 92.3%, $P=.006$), mostly receiving their medical care at the VHA (471/595, 79.2% vs. 484/664, 72.9%; $P=.009$), and being encouraged by their VHA care team to use the apps (276/595, 46.4% vs 159/664, 24%; $P<.001$). In addition, veterans who self-reported using VHA health-related apps had a higher

average HCC score than those who did not (mean HCC score 1.7 vs 1.5, respectively; $P=.05$).

Factors Associated with VHA Health-Related App Use

Results from the unadjusted and adjusted multiple logistic regression models assessing factors associated with self-reported VHA health-related app use are presented in [Table 3](#). These analyses indicated that veterans who reported mostly receiving care at the VHA (OR 1.3, 95% CI 1.0-1.7), were in worse health (as assessed by HCC score; OR 1.1, 95% CI 1.0-1.2), reported owning a desktop or laptop computer (OR 1.8, 95% CI 1.1-3.1), had PTSD (OR 1.4, 95% CI 1.1-1.9), and reported having VHA health care team members encourage them to use the apps (OR 2.7, 95% CI 2.1-3.4) were more likely to self-report having used VHA health-related apps.

Table 3. Results of multiple logistic regression analysis of factors associated with Veterans Health Administration health-related app use (N=1259). * $P < .05$, ** $P < .01$, *** $P < .001$.

Variable	Unadjusted odds ratio (95% CI)	Adjusted odds ratio (95% CI)
Age older than 65 years (reference: age younger than 65 years)	1.1 (0.9-1.4)	N/A ^a
Male (reference: female)	1.0 (0.7-1.4)	N/A
Race (reference: White)		
Black	1.0 (0.7-1.4)	N/A
Other	1.2 (0.7-2.0)	N/A
Hispanic ethnicity (reference: non-Hispanic ethnicity)	1.6 (0.9 -3.0)	N/A
Relationship status: in a relationship ^b (reference: not in a relationship)	0.97 (0.8-1.2)	N/A
Education status: at least some college or vocational school (reference: less than some college education)	0.9 (0.6-1.3)	N/A
Socioeconomic status: "not very hard to pay for basics" (reference: "some hardship paying for basics")	0.8 (0.7-1.1)	N/A
Mostly receives medical care in the VHA ^c (reference: mostly receives medical care outside the VHA)	1.4** (1.1 -1.8)	1.3* (1.0 -1.7)
Travel time to VHA: >60 minutes (reference: ≤60 minutes)	0.8 (0.6 -1.0)	0.8 (0.6-1.1)
Perceived health status fair or poor (reference: good, very good, or excellent perceived health status)	1.1 (0.9-1.4)	N/A
Hierarchical Condition Community score (continuous)	1.1* (1.0-1.2)	1.1* (1.0-1.2)
Technology ownership (reference: does not own a device)		
Desktop or laptop computer	2.0* (1.2-3.3)	1.8* (1.1-3.1)
Tablet computer	1.2 (1.0-1.5)	1.2 (0.9-1.5)
Smartphone	1.4 (1.0-2.0)	1.2 (0.9-1.8)
Early technology adopter (reference: not an early technology adopter)	1.1 (0.9-1.4)	N/A
Very comfortable or confident using computers (reference: not very comfortable or confident using computers)	1.3 (0.9-1.9)	N/A
VHA health care team encouragement to use apps (reference: no health care team encouragement)	2.8*** (2.2-3.5)	2.7*** (2.1-3.4)
Health conditions^d		
Hypertension	1.0 (0.8-1.3)	N/A
Osteoarthritis	1.1 (0.9-1.4)	N/A
Diabetes	1.0 (0.8-1.2)	N/A
Depression	1.1 (0.9-1.4)	N/A
Chronic kidney disease	0.8 (0.7-1.1)	N/A
Ischemic heart disease	1.1 (0.8-1.4)	N/A
Asthma	1.0 (0.7-1.2)	N/A
Posttraumatic stress disorder	1.5** (1.1-2.0)	1.4* (1.1-1.9)
Peripheral vascular disease	1.2 (0.9-1.6)	N/A
Anxiety disorders	1.2 (0.9-1.6)	N/A
Atrial fibrillation	0.9 (0.6-1.2)	N/A
Heart failure	1.3 (0.9-1.9)	N/A
Acute myocardial infarction	0.8 (0.5-1.2)	N/A
Stroke	1.1 (0.7-1.7)	N/A
Prostate cancer	1.2 (0.7-2.0)	N/A
Traumatic brain injury	1.2 (0.7-2.0)	N/A

Variable	Unadjusted odds ratio (95% CI)	Adjusted odds ratio (95% CI)
Chronic obstructive pulmonary disease	1.0 (0.6-1.7)	N/A
Colorectal cancer	0.9 (0.4-2.2)	N/A
Lung cancer	0.8 (0.3-2.1)	N/A

^aN/A: not applicable.

^bDefined as married, in a civil union, or engaged; not being in a relationship was defined as being single, separated, divorced, or widowed.

^cVHA: Veterans Health Administration.

^dIn the prior five years.

Discussion

Principal Findings

Our analyses suggest that veterans had greater odds of self-reporting use of VHA health-related apps if they mostly received their health care from the VHA, were in worse health, owned a desktop or laptop computer, and had a PTSD diagnosis. Perhaps most importantly, our analyses also showed that health care team member encouragement to use the apps was strongly associated with self-reported use. Veterans who reported receiving encouragement from their VHA care team members had nearly 3 times higher odds of using VHA apps to manage their health than veterans who did not report receiving such encouragement.

This finding confirms results from previous surveys of veterans demonstrating a positive association between health care team member recommendations to use an app and veteran interest in app use [26]. This finding is also aligned with research that extends beyond the veteran population suggesting that health-related app adoption may be bolstered if it is recommended by a source that the target user trusts and finds credible [19], such as the target user's health care providers [14,17,33]. Taken together, our findings, along with these related studies, contribute to growing evidence regarding the importance of the role health care providers need to play to achieve widespread adoption of health-related apps. Based on this evidence, we recommend that health care systems committed to increasing the use of health-related apps in their patient populations consider how best to prepare their frontline clinical staff to engage with patients about apps that may be relevant to them. Such preparation could include, but is not limited to, educating health care team members about apps that are available for patients and evidence regarding their effectiveness, creating tools (eg, prescription pads for health-related apps or reminders and decision aids in the electronic health record) that can be used to cue action and remind patients about health care team member recommendations, and training health care team members on how best to have these conversations with patients or creating pathways for them to refer patients to other local experts who can talk with them about health-related apps.

Our analyses also revealed that certain patient demographics and health conditions were positively associated with app adoption. Prior research conducted outside the veteran population has shown that app adoption is associated with specific sociodemographic characteristics, including female sex, younger age, more education, higher socioeconomic status

(SES), and better health status [9,18]. Our results differ, however, in that we did not find associations based on age, gender, education, or SES, and found an opposite association between health status and app use, namely, we found that veterans who were in worse health had greater odds of using VHA health-related apps. This may be related to our findings on encouragement; veterans who are in poorer health and those who report mostly receiving their health care from the VHA might have more frequent interactions with their providers and the health care system, which might drive increased app use. Additional research on how patients can best integrate the health-related apps they adopt into their self-management practices and sustain their use over time is also needed, as are studies of how clinical team members can best integrate the data from these apps into their clinical decision-making and workflows.

In interpreting these findings, it is important to note that our sample consisted of veterans who were established health-technology users and, as such, they may have had higher levels of technology literacy. It is possible that removing technology literacy as a barrier would affect the relationship between health status and app adoption. This is particularly important because patients who have more chronic conditions may benefit more from using health-related apps and because technology literacy is a modifiable factor [34], as it can be taught. Recent systematic reviews have underscored the importance of technology literacy as a factor in the use of consumer health informatics applications [35].

Similarly, the veterans in our sample reported having learned about VHA health-related apps from a variety of sources, the most frequently reported of which were directly tied to the VHA health care system. Sources included the VHA's online patient portal, VHA health care team members, and the VHA's government website. Of note, the VHA understands the importance of its online patient portal in driving adoption of other VHA health-related apps, and leverages the system to promote, market, and direct veterans to these resources in an effort to increase engagement in care and self-management. In this way, use of one patient-facing technology can beget the use of others, suggesting the importance of interventions to support the use of other technologies (as reported by Grossman et al [36]) and their potential to indirectly impact further technology adoption. Lack of awareness or knowledge of health-related apps among patients has already been recognized as a barrier to their adoption [14,16,17,37]. Our findings suggest that, at least in the case of veterans, interactions with and resources

from the health care system might present effective opportunities for patients to learn about health-related apps and, in turn, overcome these barriers. While the VHA has demonstrated success using its online patient portal for this purpose, we recommend that health care systems also consider the potential of other patient-facing technologies available to their patient populations as potential platforms for promoting use of other health-related apps. For example, health care systems that are already using automated text-messaging systems to reach and remind patients could consider creating specific text messages designed to market the availability of other apps.

Of note, we also found that veterans with PTSD had greater odds of self-reporting use of VHA health-related apps. Interestingly, the results of our unadjusted analyses did not suggest that there were differences in app use among veterans with other diagnosed health conditions. While our data cannot speak to the specific reasons for this finding, we suspect that it may be driven in part by the fact that the VHA has more available and established apps relevant to PTSD, as well as conditions highly comorbid with PTSD, including depression, anxiety disorders, and insomnia, and these apps may thus be promoted more frequently than others.

Relatedly, a recent systematic review of available VHA and Department of Defense mental health-related apps found that while efficacy data for many such apps were emerging, research did indicate the efficacy of the PTSD Coach and Virtual Hope Box apps [23]. In addition, the VHA has several apps available to support behavioral health treatments commonly received by veterans with PTSD (ie, CPT Coach, PE Coach, and CBTi Coach), which behavioral health care providers may be encouraging veterans to use in tandem with treatment, thus bolstering adoption. Industry data also suggests that this trend is not specific to the VHA. In general, digital health products focused on psychiatric concerns have experienced more growth over the past decade than products focused on other health concerns [38]. We recommend health care systems see use of mental health-related apps as a potential opportunity to suggest other health-related apps to patients that may be valuable for addressing their other health and well-being needs.

Limitations

We cannot infer causal relationships from our analyses, and self-reported survey data are subject to biases. The veterans who compose the VET-C cohort were also intentionally sampled because they were users of another VHA patient-facing technology, the health care system's patient portal. In addition

to potentially being more likely to use technology, previous research has indicated that veterans who use the VHA's portal are more educated, younger, and have higher income than the overall veteran population [39,40], which could limit the generalizability of our findings to the overall veteran population. It is important to note, however, that to ensure the privacy and security of user's health data, many of the VHA's mobile apps require veterans to sign in through a secure sign-in partner, the options for which include a DS Logon Level 2 (Premium), ID.me, or My HealtheVet Premium account. In this way, the VET-C cohort, which comprises veterans who have a My HealtheVet Premium account, may more broadly reflect veterans who use the VHA's mobile health apps. In addition, we acknowledge that the health-related apps the VHA offers are evolving, and those available at the time we completed this project may have had differing levels of relevance to the needs of different segments of the veteran population. The VET-C cohort is also characterized by more homogeneity in important demographic factors, including education and SES levels, than the overall veteran population, which may further limit generalizability. The limited number of female veterans in our sample may have further curtailed our ability to detect differences associated with gender. Finally, as with any effort to collect longitudinal data, there was attrition between the administration of our baseline survey and our second-round survey, which could have introduced response bias.

Conclusions

In this survey of veterans, we found that nearly half of respondents self-reported use of VHA health-related apps and that encouragement from a veteran's health care team was a critical factor associated with self-reported app use. Veterans predominantly reported learning about available health-related apps through other VHA technologies or their VHA health care team members. These results add to growing evidence suggesting that endorsement of apps by a health care system or health care providers can positively impact patient uptake and use. Future work should examine approaches to supporting efforts by health care team members to engage with patients about apps that may be most beneficial to their health, as well as ways to support shared decision-making regarding which apps to use and how best to integrate them as components of care and self-management. Such approaches could be included as part of multicomponent implementation strategies and tested to determine their impacts on the adoption of health-related apps by patients.

Acknowledgments

This work was supported by the Veterans Health Administration (VHA) Office of Connected Care, VHA Office of Research and Development, VHA Health Services Research and Development Service, and VHA Quality Enhancement Research Initiative Program (PEC 15-470; principal investigator: TPH). The authors also wish to thank Angela Patterson, MA, Linda Am, MPH, and Lincoln Clarke, MSA, for their invaluable contributions to data collection and processing and their review of the data-collection instruments for readability and appropriateness.

Disclaimer

The views expressed in this article are those of the authors and do not necessarily reflect the position and/or policy of the Department of Veterans Affairs or the United States Government.

All evaluation procedures described in this manuscript were completed in accordance with the ethical standards of the responsible committee on human experimentation (institutional or regional) and with the Helsinki Declaration.

Conflicts of Interest

None declared.

References

1. Xu W, Liu Y. mHealthApps: a repository and database of mobile health apps. *JMIR Mhealth Uhealth* 2015 Mar 18;3(1):e28 [FREE Full text] [doi: [10.2196/mhealth.4026](https://doi.org/10.2196/mhealth.4026)] [Medline: [25786060](https://pubmed.ncbi.nlm.nih.gov/25786060/)]
2. Whitehead L, Seaton P. The effectiveness of self-management mobile phone and tablet apps in long-term condition management: a systematic review. *J Med Internet Res* 2016 May 16;18(5):e97 [FREE Full text] [doi: [10.2196/jmir.4883](https://doi.org/10.2196/jmir.4883)] [Medline: [27185295](https://pubmed.ncbi.nlm.nih.gov/27185295/)]
3. Linardon J, Cuijpers P, Carlbring P, Messer M, Fuller-Tyszkiewicz M. The efficacy of app-supported smartphone interventions for mental health problems: a meta-analysis of randomized controlled trials. *World Psychiatry* 2019 Oct;18(3):325-336 [FREE Full text] [doi: [10.1002/wps.20673](https://doi.org/10.1002/wps.20673)] [Medline: [31496095](https://pubmed.ncbi.nlm.nih.gov/31496095/)]
4. Lecomte T, Potvin S, Corbière M, Guay S, Samson C, Cloutier B, et al. Mobile apps for mental health issues: meta-review of meta-analyses. *JMIR Mhealth Uhealth* 2020 May 29;8(5):e17458 [FREE Full text] [doi: [10.2196/17458](https://doi.org/10.2196/17458)] [Medline: [32348289](https://pubmed.ncbi.nlm.nih.gov/32348289/)]
5. Zhao J, Freeman B, Li M. Can mobile phone apps influence people's health behavior change? an evidence review. *J Med Internet Res* 2016 Oct 31;18(11):e287 [FREE Full text] [doi: [10.2196/jmir.5692](https://doi.org/10.2196/jmir.5692)] [Medline: [27806926](https://pubmed.ncbi.nlm.nih.gov/27806926/)]
6. Pérez-Jover V, Sala-González M, Guilabert M, Mira JJ. Mobile apps for increasing treatment adherence: systematic review. *J Med Internet Res* 2019 Jun 18;21(6):e12505 [FREE Full text] [doi: [10.2196/12505](https://doi.org/10.2196/12505)] [Medline: [31215517](https://pubmed.ncbi.nlm.nih.gov/31215517/)]
7. Krebs P, Duncan DT. Health app use among US mobile phone owners: a national survey. *JMIR Mhealth Uhealth* 2015 Nov 04;3(4):e101 [FREE Full text] [doi: [10.2196/mhealth.4924](https://doi.org/10.2196/mhealth.4924)] [Medline: [26537656](https://pubmed.ncbi.nlm.nih.gov/26537656/)]
8. Ramirez V, Johnson E, Gonzalez C, Ramirez V, Rubino B, Rossetti G. Assessing the use of mobile health technology by patients: an observational study in primary care clinics. *JMIR Mhealth Uhealth* 2016 Apr 19;4(2):e41 [FREE Full text] [doi: [10.2196/mhealth.4928](https://doi.org/10.2196/mhealth.4928)] [Medline: [27095507](https://pubmed.ncbi.nlm.nih.gov/27095507/)]
9. Robbins R, Krebs P, Jagannathan R, Jean-Louis G, Duncan DT. Health app use among US mobile phone users: analysis of trends by chronic disease status. *JMIR Mhealth Uhealth* 2017 Dec 19;5(12):e197 [FREE Full text] [doi: [10.2196/mhealth.7832](https://doi.org/10.2196/mhealth.7832)] [Medline: [29258981](https://pubmed.ncbi.nlm.nih.gov/29258981/)]
10. Tarricone R, Cucciniello M, Armeni P, Petracca F, Desouza KC, Hall LK, et al. Mobile health divide between clinicians and patients in cancer care: results from a cross-sectional international survey. *JMIR Mhealth Uhealth* 2019 Sep 06;7(9):e13584 [FREE Full text] [doi: [10.2196/13584](https://doi.org/10.2196/13584)] [Medline: [31493318](https://pubmed.ncbi.nlm.nih.gov/31493318/)]
11. Zhou L, Bao J, Watzlaf V, Parmanto B. Barriers to and facilitators of the use of mobile health apps from a security perspective: mixed-methods study. *JMIR Mhealth Uhealth* 2019 Apr 16;7(4):e11223 [FREE Full text] [doi: [10.2196/11223](https://doi.org/10.2196/11223)] [Medline: [30990458](https://pubmed.ncbi.nlm.nih.gov/30990458/)]
12. Abelson JS, Kaufman E, Symer M, Peters A, Charlson M, Yeo H. Barriers and benefits to using mobile health technology after operation: A qualitative study. *Surgery* 2017 Sep;162(3):605-611. [doi: [10.1016/j.surg.2017.05.007](https://doi.org/10.1016/j.surg.2017.05.007)] [Medline: [28651777](https://pubmed.ncbi.nlm.nih.gov/28651777/)]
13. Thornton LK, Kay-Lambkin FJ. Specific features of current and emerging mobile health apps: user views among people with and without mental health problems. *Mhealth* 2018;4:56 [FREE Full text] [doi: [10.21037/mhealth.2018.11.04](https://doi.org/10.21037/mhealth.2018.11.04)] [Medline: [30701174](https://pubmed.ncbi.nlm.nih.gov/30701174/)]
14. Jeffrey B, Bagala M, Creighton A, Leavey T, Nicholls S, Wood C, et al. Mobile phone applications and their use in the self-management of Type 2 diabetes mellitus: a qualitative study among app users and non-app users. *Diabetol Metab Syndr* 2019;11:84 [FREE Full text] [doi: [10.1186/s13098-019-0480-4](https://doi.org/10.1186/s13098-019-0480-4)] [Medline: [31636719](https://pubmed.ncbi.nlm.nih.gov/31636719/)]
15. Wildenbos GA, Jaspers MWM, Schijven MP, Dusseljee-Peute LW. Mobile health for older adult patients: Using an aging barriers framework to classify usability problems. *Int J Med Inform* 2019 Apr;124:68-77. [doi: [10.1016/j.ijmedinf.2019.01.006](https://doi.org/10.1016/j.ijmedinf.2019.01.006)] [Medline: [30784429](https://pubmed.ncbi.nlm.nih.gov/30784429/)]
16. Peng W, Kanthawala S, Yuan S, Hussain SA. A qualitative study of user perceptions of mobile health apps. *BMC Public Health* 2016 Nov 14;16(1):1158 [FREE Full text] [doi: [10.1186/s12889-016-3808-0](https://doi.org/10.1186/s12889-016-3808-0)] [Medline: [27842533](https://pubmed.ncbi.nlm.nih.gov/27842533/)]
17. Peng W, Yuan S, Holtz BE. Exploring the challenges and opportunities of health mobile apps for individuals with type 2 diabetes living in rural communities. *Telemed J E Health* 2016 Sep;22(9):733-738. [doi: [10.1089/tmj.2015.0180](https://doi.org/10.1089/tmj.2015.0180)] [Medline: [26982017](https://pubmed.ncbi.nlm.nih.gov/26982017/)]
18. Carroll JK, Moorhead A, Bond R, LeBlanc WG, Petrella RJ, Fiscella K. Who uses mobile phone health apps and does use matter? a secondary data analytics approach. *J Med Internet Res* 2017 Apr 19;19(4):e125 [FREE Full text] [doi: [10.2196/jmir.5604](https://doi.org/10.2196/jmir.5604)] [Medline: [28428170](https://pubmed.ncbi.nlm.nih.gov/28428170/)]

19. Horvath KJ, Alemu D, Danh T, Baker JV, Carrico AW. Creating effective mobile phone apps to optimize antiretroviral therapy adherence: perspectives from stimulant-using HIV-positive men who have sex with men. *JMIR Mhealth Uhealth* 2016 Apr 15;4(2):e48 [FREE Full text] [doi: [10.2196/mhealth.5287](https://doi.org/10.2196/mhealth.5287)] [Medline: [27084049](https://pubmed.ncbi.nlm.nih.gov/27084049/)]
20. Olenick M, Flowers M, Diaz VJ. US veterans and their unique issues: enhancing health care professional awareness. *Adv Med Educ Pract* 2015;6:635-639 [FREE Full text] [doi: [10.2147/AMEP.S89479](https://doi.org/10.2147/AMEP.S89479)] [Medline: [26664252](https://pubmed.ncbi.nlm.nih.gov/26664252/)]
21. Eibner C, Krull H, Brown K, Cefalu M, Mulcahy AW, Pollard M, et al. Current and projected characteristics and unique health care needs of the patient population served by the Department of Veterans Affairs. *Rand Health Q* 2016 May 09;5(4):13 [FREE Full text] [Medline: [28083423](https://pubmed.ncbi.nlm.nih.gov/28083423/)]
22. Zulman DM, Pal Chee C, Wagner TH, Yoon J, Cohen DM, Holmes TH, et al. Multimorbidity and healthcare utilisation among high-cost patients in the US Veterans Affairs Health Care System. *BMJ Open* 2015 Apr 16;5(4):e007771 [FREE Full text] [doi: [10.1136/bmjopen-2015-007771](https://doi.org/10.1136/bmjopen-2015-007771)] [Medline: [25882486](https://pubmed.ncbi.nlm.nih.gov/25882486/)]
23. Gould CE, Kok BC, Ma VK, Zapata AML, Owen JE, Kuhn E. Veterans Affairs and the Department of Defense mental health apps: A systematic literature review. *Psychol Serv* 2019 May;16(2):196-207. [doi: [10.1037/ser0000289](https://doi.org/10.1037/ser0000289)] [Medline: [30431306](https://pubmed.ncbi.nlm.nih.gov/30431306/)]
24. Bush NE, Smolenski DJ, Denneson LM, Williams HB, Thomas EK, Dobscha SK. A virtual hope box: randomized controlled trial of a smartphone app for emotional regulation and coping with distress. *Psychiatr Serv* 2017 Apr 01;68(4):330-336. [doi: [10.1176/appi.ps.201600283](https://doi.org/10.1176/appi.ps.201600283)] [Medline: [27842473](https://pubmed.ncbi.nlm.nih.gov/27842473/)]
25. Possemato K, Kuhn E, Johnson E, Hoffman JE, Owen JE, Kanuri N, et al. Using PTSD Coach in primary care with and without clinician support: a pilot randomized controlled trial. *Gen Hosp Psychiatry* 2016;38:94-98. [doi: [10.1016/j.genhosppsych.2015.09.005](https://doi.org/10.1016/j.genhosppsych.2015.09.005)] [Medline: [26589765](https://pubmed.ncbi.nlm.nih.gov/26589765/)]
26. Lipschitz J, Miller CJ, Hogan TP, Burdick KE, Lippin-Foster R, Simon SR, et al. Adoption of mobile apps for depression and anxiety: cross-sectional survey study on patient interest and barriers to engagement. *JMIR Ment Health* 2019 Jan 25;6(1):e11334 [FREE Full text] [doi: [10.2196/11334](https://doi.org/10.2196/11334)] [Medline: [30681968](https://pubmed.ncbi.nlm.nih.gov/30681968/)]
27. Erbes CR, Stinson R, Kuhn E, Polusny M, Urban J, Hoffman J, et al. Access, utilization, and interest in mHealth applications among veterans receiving outpatient care for PTSD. *Mil Med* 2014 Nov;179(11):1218-1222. [doi: [10.7205/MILMED-D-14-00014](https://doi.org/10.7205/MILMED-D-14-00014)] [Medline: [25373044](https://pubmed.ncbi.nlm.nih.gov/25373044/)]
28. Etingen B, Amante DJ, Martinez RN, Smith BM, Shimada SL, Richardson L, et al. Supporting the implementation of connected care technologies in the veterans health administration: cross-sectional survey findings from the veterans engagement with technology collaborative (vet-c) cohort in the Veterans Health Administration: cross-sectional survey findings from the Veterans Engagement with Technology Collaborative (VET-C) Cohort. *J Particip Med* 2020 Sep 30;12(3):e21214 [FREE Full text] [doi: [10.2196/21214](https://doi.org/10.2196/21214)] [Medline: [33044944](https://pubmed.ncbi.nlm.nih.gov/33044944/)]
29. 36-Item Short Form Survey (SF-36). RAND Corporation. URL: https://www.rand.org/health-care/surveys_tools/mos/36-item-short-form.html [accessed 2022-12-01]
30. Cho AH, Arar NH, Edelman DE, Hartwell PH, Oddone EZ, Yancy WS. Do diabetic veterans use the Internet? Self-reported usage, skills, and interest in using My HealthVet Web portal. *Telemed J E Health* 2010 Jun;16(5):595-602. [doi: [10.1089/tmj.2009.0164](https://doi.org/10.1089/tmj.2009.0164)] [Medline: [20575727](https://pubmed.ncbi.nlm.nih.gov/20575727/)]
31. Pope GC, Kautter J, Ellis RP, Ash AS, Ayanian JZ, Lezzoni LI, et al. Risk adjustment of Medicare capitation payments using the CMS-HCC model. *Health Care Financ Rev* 2004;25(4):119-141 [FREE Full text] [Medline: [15493448](https://pubmed.ncbi.nlm.nih.gov/15493448/)]
32. Morgan RO, Petersen LA, Hasche JC, Davila JA, Byrne MM, Osemene NI, et al. VHA pharmacy use in veterans with Medicare drug coverage. *Am J Manag Care* 2009 Mar 16;15(3):e1-e8 [FREE Full text] [Medline: [19298095](https://pubmed.ncbi.nlm.nih.gov/19298095/)]
33. Jacomet C, Ologeanu-Taddei R, Prouteau J, Lambert C, Linard F, Bastiani P, et al. E-health. Patterns of use and perceived benefits and barriers among people living with HIV and their physicians. Part 2: Health apps and smart devices. *Med Mal Infect* 2020 Oct;50(7):582-589. [doi: [10.1016/j.medmal.2020.04.005](https://doi.org/10.1016/j.medmal.2020.04.005)] [Medline: [32302672](https://pubmed.ncbi.nlm.nih.gov/32302672/)]
34. Alkureishi MA, Choo Z, Rahman A, Ho K, Benning-Shorb J, Lenti G, et al. Digitally disconnected: qualitative study of patient perspectives on the digital divide and potential solutions. *JMIR Hum Factors* 2021 Dec 15;8(4):e33364 [FREE Full text] [doi: [10.2196/33364](https://doi.org/10.2196/33364)] [Medline: [34705664](https://pubmed.ncbi.nlm.nih.gov/34705664/)]
35. Gibbons MC, Wilson RF, Samal L, Lehmann CU, Dickersin K, Lehmann HP, et al. Consumer health informatics: results of a systematic evidence review and evidence based recommendations. *Transl Behav Med* 2011 Mar;1(1):72-82 [FREE Full text] [doi: [10.1007/s13142-011-0016-4](https://doi.org/10.1007/s13142-011-0016-4)] [Medline: [24073033](https://pubmed.ncbi.nlm.nih.gov/24073033/)]
36. Grossman LV, Masterson Creber RM, Benda NC, Wright D, Vawdrey DK, Ancker JS. Interventions to increase patient portal use in vulnerable populations: a systematic review. *J Am Med Inform Assoc* 2019 Aug 01;26(8-9):855-870 [FREE Full text] [doi: [10.1093/jamia/ocz023](https://doi.org/10.1093/jamia/ocz023)] [Medline: [30958532](https://pubmed.ncbi.nlm.nih.gov/30958532/)]
37. Schueller SM, Neary M, O'Loughlin K, Adkins EC. Discovery of and interest in health apps among those with mental health needs: survey and focus group study. *J Med Internet Res* 2018 Jun 11;20(6):e10141 [FREE Full text] [doi: [10.2196/10141](https://doi.org/10.2196/10141)] [Medline: [29891468](https://pubmed.ncbi.nlm.nih.gov/29891468/)]
38. McCarthy O. Looking Back and Ahead at the Digital Therapeutics Industry. *MedRhythms*. 2020. URL: <https://medrhythms.medium.com/looking-back-ahead-at-the-digital-therapeutics-industry-2e8454d342c5> [accessed 2022-12-01]

39. Shimada SL, Allison JJ, Rosen AK, Feng H, Houston TK. Sustained use of patient portal features and improvements in diabetes physiological measures. *J Med Internet Res* 2016 Jul 01;18(7):e179 [FREE Full text] [doi: [10.2196/jmir.5663](https://doi.org/10.2196/jmir.5663)] [Medline: [27369696](https://pubmed.ncbi.nlm.nih.gov/27369696/)]
40. Shimada SL, Brandt CA, Feng H, McInnes DK, Rao SR, Rothendler JA, et al. Personal health record reach in the Veterans Health Administration: a cross-sectional analysis. *J Med Internet Res* 2014 Dec 12;16(12):e272 [FREE Full text] [doi: [10.2196/jmir.3751](https://doi.org/10.2196/jmir.3751)] [Medline: [25498515](https://pubmed.ncbi.nlm.nih.gov/25498515/)]

Abbreviations

CDW: corporate data warehouse
HCC: Hierarchical Condition Community
OR: odds ratio
PTSD: posttraumatic stress disorder
SES: socioeconomic status
VET-C: Veterans Engagement with Technology Collaborative
VHA: Veterans Health Administration

Edited by L Buis; submitted 09.08.22; peer-reviewed by R Marshall; comments to author 01.09.22; revised version received 26.10.22; accepted 22.11.22; published 30.12.22

Please cite as:

Hogan TP, Etingen B, Lipschitz JM, Shimada SL, McMahon N, Bolivar D, Bixler FR, Irvin D, Wacks R, Cutrona S, Frisbee KL, Smith BM

Factors Associated With Self-reported Use of Web and Mobile Health Apps Among US Military Veterans: Cross-sectional Survey
JMIR Mhealth Uhealth 2022;10(12):e41767

URL: <https://mhealth.jmir.org/2022/12/e41767>

doi: [10.2196/41767](https://doi.org/10.2196/41767)

PMID:

©Timothy P Hogan, Bella Etingen, Jessica M Lipschitz, Stephanie L Shimada, Nicholas McMahon, Derek Bolivar, Felicia R Bixler, Dawn Irvin, Rachel Wacks, Sarah Cutrona, Kathleen L Frisbee, Bridget M Smith. Originally published in JMIR mHealth and uHealth (<https://mhealth.jmir.org>), 30.12.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR mHealth and uHealth, is properly cited. The complete bibliographic information, a link to the original publication on <https://mhealth.jmir.org/>, as well as this copyright and license information must be included.